



# Clinical Practice & Epidemiology in Mental Health

Content list available at: <https://clinical-practice-and-epidemiology-in-mental-health.com>



## REVIEW ARTICLE

# Typology of Social Network Structures and Late-Life Depression in Low- and Middle-Income Countries

Akin Ojagbemi<sup>1,\*</sup> and Oye Gureje<sup>1</sup>

<sup>1</sup>World Health Organization (WHO) Collaborating Centre for Research and Training in Mental Health, Neurosciences, and Substance Abuse, Department of Psychiatry, University of Ibadan, Ibadan, Nigeria

### Abstract:

#### Background:

Rapid social changes and youth migration ensures a continuous drain on the social networks of the elderly in Low- and Middle-Income Countries (LMICs).

#### Objective:

We reviewed available literature on the relationship between social network structures and depression among community dwelling older persons in LMICs with a view to identifying patterns that might provide information for designing preventive psychosocial interventions.

#### Methods:

We searched the MEDLINE database through Pubmed, extracted information on the typologies of social network structures in LMICs and identified dimensions with the strongest systematic association with late-life depression, by weight, using the inverse of variance method. All analyses were conducted using the Cochrane review manager version 5.3.

#### Results:

Fourteen community-based surveys drawn from 16 LMIC contexts met criteria for syntheses. They included a total of 37,917 mostly female (58.8%) participants with an average age of 73.2 years. Social network size, contact with network, diversity of network, co-residency with own child, having more friends than family in the network, and prestigious standing of persons in the social network were protective structures against late-life depression. Conversely, low network diversity contributed 44.2% of the weight of all social network structures that are predictive of late-life depression.

#### Conclusion:

Recommendations are made for the design of new measures of social network structures in LMICs that captures the key dimensions identified. Epidemiological studies using such tools will provide more precise information for planning and prioritization of scarce resources for the prevention of late-life depression in LMICs.

**Keywords:** Social relationships, Social support, Socio-economic transition, Extended family systems, Socio-economic context, Socio-cultural context.

### Article History

Received: July 25, 2019

Revised: October 08, 2019

Accepted: October 10, 2019

## 1. INTRODUCTION

By 2050, approximately 80% of older persons in the world will be residents of Low and Middle Income Countries (LMICs) [1 - 4]; depression is the most common mental health

condition in old age [5, 6]. Some observational studies suggest that the prevalence and incidence of late-life depression may be much higher in LMICs than those reported for high income countries [7]. Consequently, research has focused on possible preventive factors for late-life depression. In particular, the observation that late-life depression may have a strong relationship with contextual factors in the social environment of older people [8] has propelled hypotheses around the

\* Address correspondence to this author at the Department of Psychiatry, College of Medicine, University of Ibadan, P.M.B 5017 (G.P.O) Ibadan, Nigeria; Tel: +234-8036737171; E-mails: [drakinjagbemi@yahoo.com](mailto:drakinjagbemi@yahoo.com); [akinojagbemi@gmail.com](mailto:akinojagbemi@gmail.com)

possible protective effect of good social network [9].

Social networks are the systems of social relationships in which an individual is embedded [10]. As people age and experience many losses, their social networks also narrow. This, in turn, reduces the amount of social resources available to cope with the accumulation of stressors in old age [11, 12]. In this way, persons with limited social network are likely to be at increased risk of late-life depression. In reference to another frequently cited theory [13], social network is also seen to be influenced by culture, politics and economic factors.

In terms of structure, a person's social network may consist of people with whom they regularly interact, the characteristics of such individuals, and the quality of interactions [9]. Social networks have also functional characteristics by increasing the social support resources available to the individual [14]. Yet, the availability of a social network structure does not necessarily translate to function [15]. Therefore, the two components can be studied independently.

Many LMICs are undergoing rapid socio-economic transition [16]. Poverty, material deprivation, armed conflicts, and general insecurity are widespread [17, 18]. Health and social care is often unavailable for the majority of older people in the region [19]. Moreover, youth migration is occurring faster than ever before [20, 21]. Rapid youth migration, in particular, ensures a continuous drain on the social networks of older people through the erosion traditional of extended family systems [22, 23].

The unique socio-economic and cultural contexts of LMICs suggest that the conceptualisation of social network structures is likely to be different from those of high-income countries. Few studies have considered the typologies of social network structures and how these might relate to prevalence, onset and course of late-life depression. The objective of the present study was to review the literature on the relationship between social network structures and depression among community dwelling elderly persons in LMICs and identify the most important protective typologies.

## 2. METHODS

### 2.1. Search Strategy

We searched the Medline database through PUBMED on the 13<sup>th</sup> of May 2019 using the following keywords: 'depression OR depressive AND Social network OR Networks OR Networking'. Searches were limited to English language literature and those based on human samples who were aged 60 years or older.

### 2.2. Inclusion Criteria

Studies were included if; 1) participants were drawn from a country grouped as belonging to LMICs in the World Health Organization and World Bank income categories [24], 2) they investigated epidemiological phenomena such as prevalence, incidence, risk or associated factors, and outcome, 3) they were conducted among community-dwelling participants, and 4)

they used epidemiological or experimental study designs, such as descriptive and analytical cross-sectional studies, prospective and retrospective cohort studies, case control studies, randomized controlled trials, non-randomized controlled trials, quasi-experimental, as well as before and after studies.

For the purpose of inclusion, social network structure was defined as the organization (rather than its functions) of an older person's social network [14]. This may include, but not limited to, a). The number of people with whom the older person regularly interacts, b). The characteristics of such individuals, c). Their relationships with the older person, and d). the quality of the interactions.

### 2.3. Exclusion Criteria

We excluded the following types of studies, 1) those conducted in hospitals, rehabilitation settings, nursing homes or other such institutions, 2) review papers, case series, individual case reports, expert opinions, discussion papers, and position papers; and 3) studies focusing solely on qualitative data. Study assessment for inclusion and exclusion criteria as well as subsequent data extraction was conducted based on the descriptions in the original article.


### 2.4. Statistical Methods

Flexible random effect meta-analyses were conducted to investigate the most important social network typologies that are protective or predictive of late-life depression. For these analyses, we included only studies reporting statistically significant effects at below 5% alpha. We then entered the log of study effect sizes as well as their corresponding Standard Errors (S.E) into the meta-analytic models. Estimates of effect sizes and S.Es were derived from data in the original article using methodologies developed by the Cochrane collaboration [25]. Weighting was applied using the inverse of variance method.

To determine the extent of statistical heterogeneity, we estimated the percentage of total variation in estimates reported across studies that is due to heterogeneity, rather than chance. This was computed using the  $I^2$  test. All analyses were conducted using the Cochrane review manager (Revman) version 5.3 software [26].

## 3. RESULTS

The combined database searches identified a total of 890 records. From an initial screening of the titles and abstracts of these records, we identified 24 articles that might have contained information relevant to the review. After retrieving and reading through the full texts of all 24 articles, a further 10 articles were excluded for reasons presented in Fig. (1): four studies [2, 27 - 29] investigated constructs other than social network structures, three studies [3, 4, 30] were based on general, rather than elderly, population samples (and did not present results specific to older participants in their samples), two studies [31, 32] were conducted among high income populations, and one study [33] was reported in Spanish.



**Fig. (1).** Preferred reporting items for systematic reviews and meta-analyses flow chart for the study of typologies of social network structures and late-life depression in low- and middle-income countries.

### 3.1. Appraisal of Studies Meeting Inclusion Criteria

In all, 14 studies met criteria for syntheses. The studies

were drawn from across 16 LMIC contexts. They included a total of 37,917 mostly female (58.8%) participants with an average age of 73.2 years (Table 1).

**Table 1.** Participants characteristics in studies of social network structures and late-life depression in low- and middle-income countries.

References	Low- and Middle-Income Countries	Female %	Average ages, Years
Aung <i>et al.</i> 2016	Thailand	54.9	83.0
Singh <i>et al.</i> 2016	India	53.0	65.0
Kim <i>et al.</i> 2015	Vietnam, Lao and Cambodia	49.3	67.0
Wahlin <i>et al.</i> 2015	Bangladesh	55.0	69.6
Chen <i>et al.</i> 2016	China	43.5	80.2
Monserud <i>et al.</i> 2015	Mexico	53.6	≥50 <sup>a</sup>
Gureje <i>et al.</i> 2011	Nigeria	40.4	72.4
Webster 2014	Lebanon	52	70.4
Cao, <i>et al.</i> 2015	China	62.8	60-94 <sup>b</sup>
Thiyagarajan <i>et al.</i> 2014	India, China, Cuba, Dominican Republic, Venezuela, Mexico, Peru and Puerto Rico	58.9	74
Caetano <i>et al.</i> 2013	Brazil	65.2	≥60 <sup>a</sup>
Tong <i>et al.</i> 2011	China	72.4	79.1
Chan <i>et al.</i> 2009	China	100	71
Siccote <i>et al.</i> 2008	Cuba	62.8	≥60 <sup>a</sup>

<sup>a</sup>No measure of central tendencies reported in the original article, <sup>b</sup>Only the age range was reported.

Key information about the studies is presented in Table 2. They were published between April 2008 and August 2016. About 30% of included studies were from China. Twelve (85.7%) had a cross-sectional design. The remaining two

studies were prospective observations of between two- [34] and three-years [7] follow-up. Only three studies used standardised clinical interviews for the ascertainment of depression [7, 34, 35]. The remaining 78.6% relied on various rating scales.

**Table 2. Characteristics of studies of social network structures and late-life depression in low- and middle-income countries: citations are repeated if study investigated more than one social network structure typologies.**

<i>Community Based Cross-Sectional Surveys</i>						
<i>References</i>	<i>Sample Size</i>	<i>Social Network Typology</i>	<i>Outcome Measure</i>	<i>Measure of Effect</i>	<i>Effect Size (95% C.I)<sup>a</sup> /Standard Errors</i>	<i>Effect (YES/NO)</i>
<b>Aung et al. 2016</b>	435	Size (Social Network Index)	30-item GDS <sup>b</sup>	Beta/Linear regression	Larger size, -0.18 (-0.28, -0.09)	Protective
<b>Singh et al. 2016</b>	630	Composition: children, relatives, friends, confidant networks	WHO <sup>c</sup> Composite International Diagnostic Interview	O.R <sup>d</sup> /Logistic regression	Children, 1.32 (0.79–2.21) Friends, 0.33 (0.18–0.61) Relatives, 0.70 (0.39–1.27) Confidant, 0.549 (0.17–1.75)	Protective if composed of friends
<b>Chen et al. 2016</b>	400	Composition: family, relatives, friends, health care networks	15-Item GDS <sup>b</sup>	Beta/Linear regression	Family, -0.46/0.43 Friends, -1.1/0.55 Health and social carers, 1.17/0.51	Protective if friends, Predictive if carers
<b>Kim et al. 2015</b>	147	Interaction (Lubben’s Scale)	15-item GDS <sup>b</sup>	Beta/linear regression	Greater connectedness, -0.02/0.28	NO
<b>Caetano et al. 2013</b>	3649	Interaction: Reciprocal visits	Self-report of clinician diagnoses	O.R <sup>d</sup> /Logistic regression	No visits, 1.38 (0.83 – 2.29)	NO
<b>Tong et al. 2011</b>	228	Interaction: Meetings/telephone	15-Item GDS <sup>b</sup>	Beta/Linear regression	Telephone, -0.083/8.30 Meeting with someone, -0.027/2.7	NO
<b>Wahlin et al. 2015</b>	471	Interaction: Satisfaction with contact	Self Reporting Questionnaire	Prevalence rate ratio	Fully satisfied with contact, 0.61 (0.46–0.80)	Protective
<b>Sicotte et al. 2008</b>	1905	Geographical Proximity: Co-resident with children or not	15-Item GDS <sup>b</sup>	O.R/Logistic regression	<u>Co-resident children</u> Women, 0.43, (0.25–0.74) Men, 0.29, (0.07–1.16) <u>Non co-resident children</u> Women, 0.70, (0.44–1.11) Men, 0.94, (0.40–2.20)	Protective in women with co-resident child
<b>Wahlin et al. 2015</b>	471	Geographical proximity: Co-residency	Self Reporting Questionnaire	Prevalence rate ratio	Co-resident child, 0.78 (0.66–0.94) No co-resident children, 1.28 (1.02–1.60)	Protective: co-resident Predictive: if not.
<b>Monserud et al. 2015</b>	8704	Geographical proximity: Co-residency with own children/other relative/friends/network lives in the neighborhood	Modified CES-D <sup>e</sup> 20	Beta/linear regression	Co-resident children, 0.18/0.21, Co-resident other relatives, 0.31/0.17, Children nearby, 0.05/0.10, Relatives nearby, -0.14/0.10, Friends nearby, -0.04/0.11	Predictive if co-resident with other relatives ( <i>i.e.</i> , Other than their own children)
<b>Webster 2014</b>	195	Geographical proximity: Network lives far away	CES-D <sup>e</sup> 11	Beta/linear regression	Distant family, 0.43/2.24,	NO

(Table 2) *contd.....*

<b>Webster 2014</b>	195	Emotional proximity: Hierarchical mapping	CES-D <sup>e</sup> 11	Beta/linear regression	Distant family, 2.57/1.78, Close family, -2.31/1.61	NO
<b>Cao, et al. 2015</b>	928	Social standing: Occupational Prestige Scale	30-Item GDS <sup>b</sup>	Beta/linear regression	Prestigious, -0.07/0.34	Protective
<b>Thiyagarajan, et al. 2014</b>	17,031	Diversity: PANT <sup>f</sup> assessment	Geriatric Mental State Examination	Prevalence ratio	Low diversity, 1.20 (1.08 – 1.34)	Predictive
<b>Sicotte et al. 2008</b>	1905	Diversity: Types of relationship	15-Item GDS <sup>b</sup>	O.R/Logistic regression	High diversity Women, 0.34, (0.17–0.69) Men, 0.14, (0.04–0.48)	Protective
<b>Chan et al. 2009</b>	1042	Social Network Scale (aggregate scores)	15-Item GDS <sup>b</sup>	O.R/Logistic regression	Lower social network scores, 3.63, (2.35–5.60)	Predictive
<b>COMMUNITY BASED OBSERVATIONAL STUDIES</b>						
<b>Monserud et al. 2015</b>	8704	Geographical proximity	Modified CES-D <sup>e</sup> 20	Beta/linear regression	Co-resident children, 0.20/0.23, Co-resident others, -0.31/0.18, Children nearby, -0.01/0.14, Relatives nearby, 0.25/0.13, Friends nearby, 0.24/0.13	Symptomatic improvement if co- resident with other relatives
<b>Gureje et al. 2011</b>	2149	Interaction (frequency of contacts)	WHO <sup>c</sup> Composite International Diagnostic Interview	Hazard ratios/Cox regression	<u>No Family contacts</u> women, 2.4, (0.3–20.3) men, 2.5, (0.4–16.0) <u>No friends contact</u> Women, 2.1 (1.1–3.7) Men, 1.5 (0.7–3.4)	Predictive of onset in women with no friends contact

<sup>a</sup>Final Adjusted Model, <sup>b</sup>Geriatric Depression Scale, <sup>c</sup>World Health Organization, <sup>d</sup> Odds Ratios, <sup>e</sup>Center for Epidemiological Studies Depression Scale, <sup>f</sup>Practical Assessment of Network Types.

Included studies clearly described the source population and sampling frame. Appropriate analytical techniques were used. Half of the studies derived beta values for the association between social network structures and late-life depression. The remaining studies derived other valid measures of study effect (Table 2). The corresponding S.E or 95% Confidence Intervals of the reported effect sizes were also presented.

### 3.2. Ascertainment of Social Network Structures

In Table 2, various standardised questionnaires and rating scales were used to ascertain types of social network structures. Apart from an aggregate score on the Lubben's Social Network Scale [36], six broad typologies of social network structures were investigated across studies. These include:

- [1] *Social network size*, defined in Aung *et al.* [37] as the number of persons in the social network. This was assessed using the Cohen's index [38].
- [2] *Social network composition*, defined in Singh *et al.* [39] as comprising either children network, relatives' network, friends' network or confidant network. The same construct was classified in Chen *et al.* [40] as either having family, relatives, friends or health/social care workers in the network.
- [3] *Social network interaction*, This was defined using the Lubben's Social Network Scale [36] in Kim *et al.* [41]. It was also described as the frequency of reciprocal

visits between the older persons and other people in their network [42]. Further, the interaction construct was defined in other studies as having a meeting with someone in the network [43], telephone conversation [43], satisfaction with contact [44] or regularity of contact [7].

- [4] *Geographical proximity*, This was defined by co-residency with own children (whether biological or foster) or not [34, 44, 45]. Co-residency with other family members rather than own children was an additional definition of proximity provided in Monserud and Wong [34]. For non-co-resident respondents, geographical proximity was further expounded by whether respondents had adult children who lived in the same neighborhood [34], or lived more than half an hour from the older person [46]. The emotional proximity of social network was a supplementary category of geographical proximity defined in Webster *et al.* [46] as the relative distance between the older persons and people in their network.
- [5] *Social network diversity*, Summarised in Thiyagarajan *et al.* [35] as integrated on non-integrated networks based on results of the Practical Assessment of Network Types [47]. Diversity was also defined in Sicotte *et al.* [45] by the number of different types of relationships the older person had with people in their network.

[6] *Social standing of the network*, defined in Cao *et al.* [48] as the older person’s connection with family members, friends or acquaintances in a list of 20 specific occupations with varying prestige scores. To achieve this, a position generator procedure was applied to occupational prestige scale scores [49].

### 3.3. Association of Types of Social Network Structures with Late-life Depression

In Table 2, social network interaction was the most cited broad typology of social network structures. The construct was investigated in five studies [7, 41 - 44]. However, three of the five studies of social network interaction showed no statistical effect on depression. Also notable in Table 2 are the results of

two longitudinal observations in relation to social network interaction [7] and geographical proximity [34].

The disaggregated factors in the dimensions of social network structures demonstrating systematic association with late life depression are presented in Fig. (2). In the decreasing order, the most precise protective factors were large network size, regular contact with network, high network diversity, having a co-resident child, having more friends than family in the network, and prestigious standing of persons in the social network. However, large network size and regular contact with network contributed more than one-third (33.8%) of all systematic associations with late life depression. Conversely, 44.2% of all predictive factors for late-life depression was contributed by low network diversity.




Fig. (2). Independently associated social network structures with prevalent depression in older persons from low- and middle-income countries (ranked by the inverse of variance method).

#### 4. DISCUSSION

The present study found that six broad dimensions of social network structures demonstrate significant protective effects against late-life depression in LMICs. In descending order of weight contributed, the dimensions were represented by social network size, contact with network, diversity of network, co-residency with own child, having more friends than family in the network, and prestigious standing of persons in the social network. However, large network size and regular contact with network contributed more than one-third (33.8%) of all protective structures against late life depression. Conversely, 44.2% of all social network structures that are predictive of late-life depression was contributed by low network diversity.

The finding that prestigious standing of persons in the network contributed much less (7.9%) to the pooled weight of protective structures against late-life depression may be due to the complex nature of the position generator procedure [50]. The social standing index generated by this procedure is an aggregate score comprising the number of occupations endorsed by respondents, the highest prestige score of occupations so endorsed and the range of prestige scores. Each of these components have been found to have their own unique effects on wellbeing [51]. For example, while the endorsement of higher number of occupations demonstrate strong relationship with health, prestige scores assigned to those occupations may sometimes show non-significant effects [52]. Another possible reason for an observed small effect of social standing on late-life depression is the observation that social stratifications based on occupation may be more relevant to high-income countries [53]. This is as large numbers of community-dwelling older people in LMICs engage in subsistence farming and other elementary occupations [54].

A third observation in the present review is that individual studies on the relationship between social network and late-life depression in LMICs have often selected only a few dimensions of social network structures for investigation. In this way, studies appear less comprehensive in their approach to the social network. The result of this is the observed variability in findings. Inconsistencies in the magnitude and direction of associations make it difficult to decide on the most appropriate targets for preventive interventions based on social network characteristics. Greater precision in findings will provide the information needed by policy makers to better streamline scarce resources for the prevention of late-life depression in LMICs.

It would seem that the most comprehensive assessment of the dimension of social network structures in the present review was provided by Chan *et al.* [55] since the authors relied on the summary score on the Lubben's Social Network Scale [36]. Analyzing from the dimensions of social network structures identified in the present study, the Lubben's scale demonstrates relatively superior face validity by providing some assessment of four out of the six broad dimensions identified: social network size, diversity, interaction and geographical proximity. However, the relative predictive weight of association of the Lubben's scale summary score with depression in the present study was low. While this

finding may also suggest weaknesses in the primary study [55], it provides some evidence of inadequate known group validity of the Lubben's social network scale.

We note that the validity and reliability of available measures of social networks have long been a source of concern in the literature [15, 56]. Future research should strive towards attaining the goal of finding the best measurement modalities for social network structures and functions in specific populations and disease contexts. The optimal tool for assessing social network structures relevant to late-life depression in LMICs may yet be unavailable. Such a tool should ideally produce measures that are easy to interpret and are comprehensive enough to assess the dimensions of social network structures identified in the present study.

The observations made in this limited review should be interpreted within limitations of the quality of available data. Judgments about the contributions of the different facets of the social network to late-life depression using meta-analytic weighting may be affected by methodological variability in the included studies. In this regard, it should be noted that a significant heterogeneity ( $I^2=88\%$ ,  $p<0.001$ ) was recorded when studies reporting significant protective effects of social network structures on late-life depression were combined. On a closer look at our data, the significant statistical heterogeneity found in studies reporting protective effects of the social network was due to the effect size outlier reported when examining the effect of the prestigious social standing of members of the network on late-life depression. The study by Cao and colleagues [48] conducted among 928 older residents in Hangzhou China was the only citation in the series to use the occupational prestige scale as a meaningful index of social network structure. Even though this study [48] contributed only 7.9% of the weight of all studies reporting significant protective effect of social network structures, its effect size is about half of the combined effect sizes reported by the other studies in the series.

The present mini review has other limitations. We have carried out a rapid review of the literature on social network structures and late-life depression in LMICs. As such, the search strategy has focused on records in the MEDLINE database. Records in other important databases, such as PsychInfo and Embase have not been considered. Also, manual searches of the reference list of included studies as well as grey literature were not implemented. Nevertheless, the MEDLINE is, perhaps, the largest repository of biomedical literature. To ensure that ahead of print articles are not missed, we implemented our search of the MEDLINE database through Pubmed.

#### CONCLUSION

Health care resources are limited in LMICs. As such, accurate information about the contribution of different types of social network structures that may be protective for late-life depression in these countries may be helpful in the planning and prioritization of policies and allocation of scarce resources. New measures that include the broad dimensions of social network structures identified in the present study are needed. Such new tools should be deployed in future large-scale

surveys of the mental health and wellbeing of elderly persons living in LMICs. While cross-sectional surveys are important, the use of such tools in longitudinal observations will provide even more robust information about the relative contributions of social network structures and functions to the mental health of older people in LMICs.

#### CONSENT FOR PUBLICATION

Not applicable.

#### FUNDING

None.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

#### ACKNOWLEDGEMENTS

AO performed literature search, data extraction and analyses, AO and OG wrote the first draft of the paper. Both authors approved the final draft for submission.

#### REFERENCES

- [1] Dotchin CL, Akinyemi RO, Gray WK, Walker RW. Geriatric medicine: services and training in Africa. *Age Ageing* 2013; 42(1): 124-8. [http://dx.doi.org/10.1093/ageing/afs119] [PMID: 23027519]
- [2] Dong X, Beck T, Simon MA. The associations of gender, depression and elder mistreatment in a community-dwelling Chinese population: the modifying effect of social support. *Arch Gerontol Geriatr* 2010; 50(2): 202-8. [http://dx.doi.org/10.1016/j.archger.2009.03.011] [PMID: 19398133]
- [3] Axinn WG, Ghimire DJ, Williams NE, Scott KM. Associations between the social organization of communities and psychiatric disorders in rural Asia. *Soc Psychiatry Psychiatr Epidemiol* 2015; 50(10): 1537-45. [http://dx.doi.org/10.1007/s00127-015-1042-1] [PMID: 25796491]
- [4] Kuscü MK, Dural U, Onen P, *et al.* The association between individual attachment patterns, the perceived social support, and the psychological well-being of Turkish informal caregivers. *Psychooncology* 2009; 18(9): 927-35. [http://dx.doi.org/10.1002/pon.1441] [PMID: 19140124]
- [5] Whiteford HA, Degenhardt L, Rehm J, *et al.* Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet* 2013; 382(9904): 1575-86. [http://dx.doi.org/10.1016/S0140-6736(13)61611-6] [PMID: 23993280]
- [6] Global Burden of Disease Study 2013 Collaborators, Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015 Jun; 7piiS0140-6736(15)60692
- [7] Gureje O, Oladeji B, Abiona T. Incidence and risk factors for late-life depression in the Ibadan Study of Ageing. *Psychol Med* 2011; 41(9): 1897-906. [http://dx.doi.org/10.1017/S0033291710002643] [PMID: 21275087]
- [8] Vink D, Aartsen MJ, Schoevers RA. Risk factors for anxiety and depression in the elderly: a review. *J Affect Disord* 2008; 106(1-2): 29-44. [http://dx.doi.org/10.1016/j.jad.2007.06.005] [PMID: 17707515]
- [9] Schwarzbach M, Luppá M, Forstmeier S, König HH, Riedel-Heller SG. Social relations and depression in late life—a systematic review. *Int J Geriatr Psychiatry* 2014; 29(1): 1-21. [http://dx.doi.org/10.1002/gps.3971] [PMID: 23720299]
- [10] Berkman LF, Syme SL. Social networks, host resistance, and mortality: a nine-year follow-up study of Alameda County residents. *Am J Epidemiol* 1979; 109(2): 186-204. [http://dx.doi.org/10.1093/oxfordjournals.aje.a112674] [PMID: 425958]
- [11] Thoits PA. *Sociological Approaches to Mental Illness. A Handbook for the Study of Mental Health Social Contexts, Theories, and Systems.* Cambridge, England: Cambridge University Press 2012; pp. 106-24.
- [12] Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psychol Bull* 1985; 98(2): 310-57. [http://dx.doi.org/10.1037/0033-2909.98.2.310] [PMID: 3901065]
- [13] Berkman LF, Glass T, Brissette I, Seeman TE. From social integration to health: Durkheim in the new millennium. *Soc Sci Med* 2000; 51(6): 843-57. [http://dx.doi.org/10.1016/S0277-9536(00)00065-4] [PMID: 10972429]
- [14] Sherbourne CD, Stewart AL. The MOS social support survey. *Soc Sci Med* 1991; 32(6): 705-14. [http://dx.doi.org/10.1016/0277-9536(91)90150-B] [PMID: 2035047]
- [15] O'Reilly P. Methodological issues in social support and social network research. *Soc Sci Med* 1988; 26(8): 863-73. [http://dx.doi.org/10.1016/0277-9536(88)90179-7] [PMID: 3287636]
- [16] *The International Migration Report 2017.* Geneva: United Nations 2017.
- [17] Ojagbemi A, Bello T, Luo Z, Gureje O. Living Conditions, Low Socioeconomic Position, and Mortality in the Ibadan Study of Aging. *J Gerontol B Psychol Sci Soc Sci* 2017; 72(4): 646-55. [PMID: 27038398]
- [18] Cappuccio FP, Miller MA. Cardiovascular disease and hypertension in sub-Saharan Africa: burden, risk and interventions. *Intern Emerg Med* 2016; 11(3): 299-305. [http://dx.doi.org/10.1007/s11739-016-1423-9] [PMID: 27001886]
- [19] Uwakwe R, Ibeh CC, Modebe AI, *et al.* The epidemiology of dependence in older people in Nigeria: prevalence, determinants, informal care, and health service utilization. A 10/66 dementia research group cross-sectional survey. *J Am Geriatr Soc* 2009; 57(9): 1620-7. [http://dx.doi.org/10.1111/j.1532-5415.2009.02397.x] [PMID: 19682135]
- [20] Leigh-Hunt N, Bagguley D, Bash K, *et al.* An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public Health* 2017; 152: 157-71. [http://dx.doi.org/10.1016/j.puhe.2017.07.035] [PMID: 28915435]
- [21] *Challenges and Priorities of Global Mental Health in the Sustainable Development Goals (SDG) Era.* London, United Kingdom: The Academy of Medical Sciences 2018; p. 24.
- [22] Fokkema T, De Jong Gierveld J, Dykstra PA. Cross-national differences in older adult loneliness. *J Psychol* 2012; 146(1-2): 201-28. [http://dx.doi.org/10.1080/00223980.2011.631612] [PMID: 22303621]
- [23] Dykstra PA. Older adult loneliness: myths and realities. *Eur J Ageing* 2009; 6(2): 91-100. [http://dx.doi.org/10.1007/s10433-009-0110-3] [PMID: 19517025]
- [24] *World Health Statistics.* Geneva, Switzerland: World Health Organization 2015.
- [25] *Cochrane Handbook for Systematic Reviews of Interventions Version 5.10.* Chichester, England: John Wiley 2011.
- [26] *Review Manager (RevMan) [Computer program].* The Nordic Cochrane Centre, Copenhagen: The Cochrane Collaboration 2014.
- [27] Baiyewu O, Yusuf AJ, Ogundele A. Depression in elderly people living in rural Nigeria and its association with perceived health, poverty, and social network. *Int Psychogeriatr* 2015; 27(12): 2009-15. [http://dx.doi.org/10.1017/S1041610215001088] [PMID: 26265242]
- [28] Park S, Hatim Sulaiman A, Srisurapanont M, *et al.* The association of suicide risk with negative life events and social support according to gender in Asian patients with major depressive disorder. *Psychiatry Res* 2015; 228(3): 277-82. [http://dx.doi.org/10.1016/j.psychres.2015.06.032] [PMID: 26160206]
- [29] Quattrin LB, Galli R, Moriguchi EH, Gastal FL, Pattussi MP. Collective efficacy and depressive symptoms in Brazilian elderly. *Arch Gerontol Geriatr* 2014; 59(3): 624-9. [http://dx.doi.org/10.1016/j.archger.2014.08.001] [PMID: 25183439]
- [30] Perkins JM, Nyakato VN, Kakuhikire B, *et al.* Food insecurity, social networks and symptoms of depression among men and women in rural Uganda: a cross-sectional, population-based study. *Public Health Nutr* 2018; 21(5): 838-48. [http://dx.doi.org/10.1017/S1368980017002154] [PMID: 28988551]
- [31] Chao SF. Assessing social support and depressive symptoms in older Chinese adults: a longitudinal perspective. *Aging Ment Health* 2011; 15(6): 765-74. [http://dx.doi.org/10.1080/13607863.2011.562182] [PMID: 21838514]



- [32] Chi I, Chou KL. Social support and depression among elderly Chinese people in Hong Kong. *Int J Aging Hum Dev* 2001; 52(3): 231-52. [http://dx.doi.org/10.2190/V5K8-CNMG-G2UP-37QV] [PMID: 11407488]
- [33] Gallegos-Carrillo K, Mudgal J, Sánchez-García S, *et al.* Social networks and health-related quality of life: A population based study among older adults. *Salud Publica Mex* 2009; 51(1): 6-13. [http://dx.doi.org/10.1590/S0036-36342009000100004] [PMID: 19180307]
- [34] Monsrud MA, Wong R. Depressive Symptoms Among Older Mexicans: The Role of Widowhood, Gender, and Social Integration. *Res Aging* 2015; 37(8): 856-86. [http://dx.doi.org/10.1177/0164027514568104] [PMID: 25651596]
- [35] Thiagarajan JA, Prince M, Webber M. Social support network typologies and health outcomes of older people in low and middle income countries—a 10/66 Dementia Research Group population-based study. *Int Rev Psychiatry* 2014; 26(4): 476-85. [http://dx.doi.org/10.3109/09540261.2014.925850] [PMID: 25137114]
- [36] Lubben J. Assessing social networks among elderly populations. *J Fam Med Community Health* 1988; 1988: 42-52. [http://dx.doi.org/10.1097/00003727-198811000-00008]
- [37] Aung MN, Moolphate S, Aung TN, Katonyoo C, Khamchai S, Wannakrairot P. The social network index and its relation to later-life depression among the elderly aged  $\geq 80$  years in Northern Thailand. *Clin Interv Aging* 2016; 11: 1067-74. [http://dx.doi.org/10.2147/CIA.S108974] [PMID: 27540286]
- [38] Cohen S, Doyle WJ, Skoner DP, Rabin BS, Gwaltney JM Jr. Social ties and susceptibility to the common cold. *JAMA* 1997; 277(24): 1940-4. [http://dx.doi.org/10.1001/jama.1997.03540480040036] [PMID: 9200634]
- [39] Singh L, Singh PK, Arokiasamy P. Social Network and Mental Health Among Older Adults in Rural Uttar Pradesh, India: A Cross-Sectional Study. *J Cross Cult Gerontol* 2016; 31(2): 173-92. [http://dx.doi.org/10.1007/s10823-016-9286-0] [PMID: 26879450]
- [40] Chen YY, Wong GH, Lum TY, *et al.* Neighborhood support network, perceived proximity to community facilities and depressive symptoms among low socioeconomic status Chinese elders. *Aging Ment Health* 2016; 20(4): 423-31. [http://dx.doi.org/10.1080/13607863.2015.1018867] [PMID: 25775108]
- [41] Kim T, Nguyen ET, Yuen EJ, Nguyen T, Sorn R, Nguyen GT. Differential Role of Social Connectedness in Geriatric Depression Among Southeast Asian Ethnic Groups. *Prog Community Health Partnersh* 2015; 9(4): 483-93. [http://dx.doi.org/10.1353/cpr.2015.0075] [PMID: 26639374]
- [42] Caetano SC, Silva CM, Vettore MV. Gender differences in the association of perceived social support and social network with self-rated health status among older adults: A population-based study in Brazil. *BMC Geriatr* 2013; 13: 122. [http://dx.doi.org/10.1186/1471-2318-13-122] [PMID: 24229389]
- [43] Tong HM, Lai DW, Zeng Q, Xu WY. Effects of social exclusion on depressive symptoms: elderly Chinese living alone in Shanghai, China. *J Cross Cult Gerontol* 2011; 26(4): 349-64. [http://dx.doi.org/10.1007/s10823-011-9150-1] [PMID: 21814741]
- [44] Wahlin Å, Palmer K, Sternäng O, Hamadani JD, Kabir ZN. Prevalence of depressive symptoms and suicidal thoughts among elderly persons in rural Bangladesh. *Int Psychogeriatr* 2015; 27(12): 1999-2008. [http://dx.doi.org/10.1017/S104161021500109X] [PMID: 26250141]
- [45] Sicotte M, Alvarado BE, León EM, Zunzunegui MV. Social networks and depressive symptoms among elderly women and men in Havana, Cuba. *Aging Ment Health* 2008; 12(2): 193-201. [http://dx.doi.org/10.1080/13607860701616358] [PMID: 18389399]
- [46] Webster NJ, Antonucci TC, Ajrouch KJ, Abdulrahim S. Social networks and health among older adults in Lebanon: the mediating role of support and trust. *J Gerontol B Psychol Sci Soc Sci* 2015; 70(1): 155-66. [http://dx.doi.org/10.1093/geronb/gbu149] [PMID: 25324295]
- [47] Wenger GC, Shahtahmasebi S. Survivors: Support network variation and sources of help in rural communities. *J Cross Cult Gerontol* 1991; 6(1): 41-82. [PMID: 24390432]
- [48] Cao W, Li L, Zhou X, Zhou C. Social capital and depression: evidence from urban elderly in China. *Aging Ment Health* 2015; 19(5): 418-29. [http://dx.doi.org/10.1080/13607863.2014.948805] [PMID: 25155221]
- [49] Bian Y. Source and functions of urbanites' social capital: A network approach. *Soc Sci China* 2004; 3: 136-46.
- [50] Lin N, Fu YC, Hsung RM. The Position Generator: Measurement techniques for investigations of social capital. In: *Social capital: Theory and research*. New York, NY: Aldine de Gruyter 2001.
- [51] Cheng ST, Lee CK, Chan AC, Leung EM, Lee JJ. Social network types and subjective well-being in Chinese older adults. *J Gerontol B Psychol Sci Soc Sci* 2009; 64(6): 713-22. [http://dx.doi.org/10.1093/geronb/gbp075] [PMID: 19820232]
- [52] Moore S, Daniel M, Gauvin L, Dubé L. Not all social capital is good capital. *Health Place* 2009; 15(4): 1071-7. [http://dx.doi.org/10.1016/j.healthplace.2009.05.005] [PMID: 19482506]
- [53] Grusky DB. The Past, Present, and Future of Social Inequality. *Social Stratification: Class, Race, and Gender in Sociological Perspective*. Boulder, CO: Westview Press 2001; pp. 3-51.
- [54] Ojagbemi A, Bello T, Gureje O. Gender differential in social and economic predictors of incident major depressive disorder in the Ibadan Study of Ageing. *Soc Psychiatry Psychiatr Epidemiol* 2018; 53(4): 351-61. [http://dx.doi.org/10.1007/s00127-018-1500-7] [PMID: 29468523]
- [55] Chan MF, Zeng W. Investigating factors associated with depression of older women in Macau. *J Clin Nurs* 2009; 18(21): 2969-77. [http://dx.doi.org/10.1111/j.1365-2702.2009.02867.x] [PMID: 19732244]
- [56] Butts CT. Social network analysis: A methodological introduction. *Asian J Soc Psychol* 2008; 11: 13-41. [http://dx.doi.org/10.1111/j.1467-839X.2007.00241.x]