DOI: 10.2174/0117450179293069250507074009, 2025, 21, e17450179293069

RESEARCH ARTICLE

Effects of a Playoff Match on Competitive Anxiety and Autonomic Regulation in Professional esports Players

Sergio Machado^{*, 1,2,3}, Leandro de Oliveira Sant'Ana⁴, Luis Cid^{5,6,7}, Diogo S. Teixeira^{8,9}, Antonio Egidio Nardi¹, Bruno Travassos^{2,5,11}, and Diogo Monteiro^{5,7,10}

¹ Laboratory of Panic and Respiration, Institute of Psychiatry (IPUB), Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

²Department of Sports Science, University of Beira Interior, 6201-001 Covilhã, Portugal

³Laboratory of Physical Activity Neuroscience, Neurodiversity Institute, Queimados 26325-020, Brazil

⁴Postgraduate Program in Physical Education, Federal University of Juiz de Fora, Juiz de Fora, Brazil

⁵Research Center in Sport, Health and Human Development (CIDESD), 5000-558 Vila Real, Portugal

⁶Sport Sciences School of Rio Maior, Polytechnic of Santarém (ESDRM-IP Santarém), Santarém, Portugal

⁷Life Quality Research Centre (CIEQV), 2400-901 Leiria, Portugal

⁸Faculty of Physical Education and Sport, Lusófona University, 1749-024 Lisbon, Portugal

[°]Research Center in Sport, Physical Education, and Exercise and Health (CIDEFES), (CIDEFES), 1749-024 Lisbon, Portugal

¹⁰ESECS, Polytechnic of Leiria, 2411-901 Leiria, Portugal

¹¹Portugal Football School, Portuguese Football Federation, 1495-433 Cruz Quebrada, Portugal

Abstract:

Introduction: A competition is considered a stressful situation since it causes physiological and emotional changes in the responses of athletes and consequently influences their performance. The aim of our study was to investigate competitive anxiety and heart rate variability (HRV) in professional eSports athletes, comparing the responses before and after matches based on whether they won or lost. We hypothesized that victorious players would display more favorable autonomic and anxiety-related responses after the matches compared to those who were defeated.

Methods: We recruited fifty male esports players from 10 different Brazilian teams and carried out the experiment across two sessions. Initially, 24 hours before the game, players signed the informed consent form, and sample characterization, along with player familiarization with anxiety and HRV, was performed. Following this, the players recorded their anxiety levels and HRV at rest for 10 minutes, both 60 and 30 minutes prior to the game (baseline time points), as well as 10 minutes after the conclusion of the game.

Results: Regarding anxiety, our results demonstrated that the victory group (VG) exhibited significantly lower scores for both cognitive and somatic anxiety in the post-game time point, coupled with increased scores for self-confidence when compared to the baseline (BL) and pre-game time points. In opposition, the defeated group (DG) displayed significantly elevated scores for cognitive and somatic anxiety during the post-game time point, accompanied by decreased self-confidence scores compared to the baseline and pre-game time points. Regarding heart rate variability (HRV), the victory group (VG) demonstrated a significant increase in SDNN, rMSSD, and HF measures, coupled with a significant decrease in the LF/HF ratio. Conversely, the defeated group (DG) exhibited a significant decrease in SDNN and rMSSD, along with a significant increase in the LF/HF ratio.

Discussion: Our results revealed that VG exhibited better HRV responses, indicating greater parasympathetic activation. VG also showed lower levels of cognitive and somatic anxiety and higher levels of self-confidence in the post-game time. In contrast, DG demonstrated worse HRV responses, indicating greater sympathetic activation, along with higher levels of cognitive and somatic anxiety and lower levels of self-confidence in the same post-game period.

Conclusion: In summary, the VG exhibited superior HRV responses in conjunction with lower levels of anxiety compared to the DG.

Keywords: Victory, Defeat, Esports, Competitive anxiety, Heart rate variability, HRV.

OPEN ACCESS



© 2025 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: https://creativecommons.org/licenses/by/4.0/legalcode. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

* Address correspondence to this author at the Laboratory of Panic and Respiration, Institute of Psychiatry (IPUB), Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil, Department of Sports Science, University of Beira Interior, 6201-001Covilhã, Portugal and Laboratory of Physical Activity Neuroscience, Neurodiversity Institute, Queimados 26325-020, Brazil; E-mail: secm80@gmail.com

Cite as: Machado S, Sant'Ana L, Cid L, Teixeira D, Nardi A, Travassos B, Monteiro D. Effects of a Playoff Match on Competitive Anxiety and Autonomic Regulation in Professional esports Players. Clin Pract Epidemiol Ment Health, 2025; 21: e17450179293069. http://dx.doi.org/10.2174/0117450179293069250507074009



Received: July 29, 2024 Revised: January 26, 2025 Accepted: January 29, 2025 Published: May 22, 2025



Send Orders for Reprints to reprints@benthamscience.net

1. INTRODUCTION

Competition is considered a stressful situation since it causes physiological and emotional changes, such as anxiety, in the responses of the athletes and consequently influences their performance [1]. Anxiety is inherent to sports, as it can be experienced at different levels [2]. A higher state of anxiety that occurs before or during sports competitions is defined as competitive anxiety [3], which can be classified as cognitive and somatic anxiety [4].

Cognitive anxiety is characterized by negative expectations about performance success or selfevaluation, accompanied by adverse feelings, thoughts, fears, and worries regarding one's performance. This can lead to a decline in self-esteem and self-confidence, negative self-talk, difficulty concentrating, and disrupted attention. In contrast, somatic anxiety is characterized by heightened physiological arousal, including increased heart rate and muscle tension. This can result in negative symptoms, such as nervousness, breathing difficulties, elevated blood pressure, and muscle tightness [4]. These changes in emotional states can occur before, during, or after sports competitions [2]. Thus, an increase in anxiety can lead to negative consequences in the athletes' performance [5, 6]. There is a scarcity of studies on esports competition as a whole, and the impact of a match physiological and psychological variables on in professional esports players is of great interest to researchers and coaches [7, 8]. As the sport continues to evolve rapidly, gaining insights into its psychophysiological aspects will be crucial for informing technical, tactical, and training-related adjustments.

Esports, better known as electronic sports, are competitions involving video games, e.g., League of Legends, and their participants, referred to as players, may compete at amateur or professional levels [8]. From a psychophysiological point of view, the autonomic nervous system (ANS) shows changes in sympathetic and parasympathetic activities [9]. Considering the sports context, the ANS assessment can be performed noninvasively through heart rate variability (HRV) [10]. In fact, HRV is a sensitive physiological indicator for psychological regulation [11] that detects changes in sympathetic/parasympathetic balance stressful in situations, such as sports competitions [12].

Research carried out by Mateo *et al.* [13] and Bernston *et al.* [14] reported that HRV is a sensitive indicator of

changes in emotional state, thus suggesting an interaction between cardiac and neural processes. It is well known that cardiac control and response to different stressors can be explained by both variations in activation of the sympathetic nervous system (SNS) [15] and parasympathetic nervous system (PNS) [16-18]. In line with this, there is abundant evidence showing that a decrease in HRV parameters is associated with anxiety [19-21]. For example, Friedman [16] and Cohen and Benjamin [22] revealed that reductions in HRV parameters correspond to an increase in anxiety. These findings prompt us to explore the impacts of match outcomes on competitive anxiety and heart rate variability (HRV). This is important, as esports appear to rely much on psychological and neurocognitive abilities for success compared to traditional sports [8]. From a practical perspective, some studies addressed competitive anxiety [23-26] and HRV responses [27] to successful or unsuccessful match outcomes in traditional sports. However, there are no studies addressing competitive anxiety and HRV responses to successful or unsuccessful match outcomes (i.e., victory and defeat) in professional esports players. A few similar studies were conducted by Pena [27] and Fuentes-Garcia et al. [28], both on tennis players. Pena [27] investigated HRV before and after competition in junior tennis players. Results indicated that losers showed a reduction in PNN50 both on the training day and after the match compared with winners. A study by Fuentes-Garcia et al. [28] examined the influence of match results (win or loss) in an international competition on pre-competition anxiety and self-assurance among elite junior tennis players. The research revealed substantial differences in cognitive anxiety and self-confidence between victors and defeated athletes when comparing their pre-match and post-match values. More specifically, the post-match evaluations of the victorious athletes revealed a reduction in cognitive anxiety and an increase in self-confidence. Regarding esports, the existing studies mainly deal with reports of playing video games or have analyzed the lifestyles and health-related issues of the players. Thus, there is a need to understand the variations in cardiovascular, respiratory, metabolic [29], and psychological health [30] in esports in different contexts or according to different game results. The existing academic literature lacks any published research that specifically examines the impacts of both victory and defeat on competitive anxiety and heart rate variability

(HRV) among professional esports athletes. A better understanding of competitive anxiety and HRV behaviors could contribute to enhancing athletic performance [31-34], thus allowing for the assessment of the recovery process between matches and throughout the competition [35].

The objective of this study was to analyze the impact of the outcome of a playoff game (victory and defeat) on the levels of competitive anxiety and HRV in professional esports athletes. More precisely, the study tested the following hypothesis: victorious players will exhibit lower levels of both cognitive and somatic anxiety, higher selfconfidence, and superior autonomic responses compared to defeated athletes in the aftermath of the match.

2. METHODS

2.1. Participants

The required sample size for this study was calculated using the G*Power v.3.1 software [36]. The input parameters used in the power analysis were a medium anticipated effect size for a comparison between two dependent means (d = 0.50), a statistical power (1- β) set at 0.80, and a significance level (α) of 0.05. Since no previous studies had provided results directly relevant to the targeted effect in our research, the researchers instead used effect size estimates from similar experimental manipulations involving psychophysiological responses in the context of competitive stress within sports science [37]. Notably, the effect sizes observed in the referenced study were found to be large [37].

The target sample size determined for our study was 20 participants $(1-\beta = 1.616)$, supported by power calculations. Fifty male esports players between the ages of 18 and 29 years (mean age: 24.98 ± 2.59 years, height: 178.6 ± 1.45 cm, weight: 78.5 ± 2.35 kg, experience duration: 7.68 ± 1.33 years, and physically inactive for 36 ± 1.05 minutes per week) from several Brazilian teams took part in the study. To be eligible, participants had to be professional athletes with at least 5 years of experience in national or international competitions and engaged in 8 to 10 hours of daily training. Exclusion criteria included having neuropsychiatric, cardiovascular, or osteoarticular diseases, using neuropsychiatric drugs, and consuming caffeinated or alcoholic beverages on the day of the experiment. The study was approved by the Ethics Committee of the University of Beira Interior (D2574), and all athletes were instructed about the risks and benefits related to the study before providing written informed consent.

2.2. Experimental Design

Participants were recruited from 10 distinct Brazilian teams, each comprising 5 players. If any individual failed to satisfy the inclusion criteria, the entire team was omitted from the study. The research methodology entailed conducting the study with two teams per pivotal game, enabling equal observation of the phenomena of triumph and defeat. As a result, upon completing the experiment, there were 25 players in the victory group (VG) and defeated group (DG). Players were exposed to pivotal games in international tournaments, such as the CS: GO PGL Major Championship Fall and Six Invitational. The study unfolded across two sessions.

In the first session, 24 hours before the game, players signed the informed consent form in the game house. Participant characteristics, such as anthropometric data (age, stature, and body mass), physical activity levels evaluated through the International Physical Activity Questionnaire (IPAQ) [38], and expertise time, were assessed for sample description. Furthermore, the Revised Competitive State Anxiety Inventory 2 (CSAI-2R) and heart rate variability (HRV) assessments were carried out in individuals.

During the second session, participants completed the CSAI-2R [39] and had their heart rate variability (HRV) recorded at rest for 10 minutes, 60 minutes, and 30 minutes prior to the match (baseline [BL] and pre-game time points, respectively) as well as 10 minutes after the conclusion of the match (post-game time point, Fig. 1). Both sessions took place at the gaming residence between 14:00 - 17:00 to mitigate the influence of circadian rhythms on psychological and autonomic function. An investigator evaluated the psychological variables and HRV.

2.3. Familiarization

The familiarization process with the CSAI-2R was conducted by the same investigator 24 hours before the experimental conditions, and it involved the following steps: 1) the researcher provided a thorough reading of the detailed instructions for each query; 2) It was clarified that responses should be selected from the options offered and that there are no right or wrong answers; 3) players were told not to answer the same question again, stressing the value of being truthful in their comments; 4) Prior to finishing the questionnaire, players were instructed to review their responses.

The familiarization process took place only prior to the experimental phase. The players then individually completed the questionnaire during the experimental conditions. Players were instructed to use the heart rate monitor, position the chest strap over the xiphoid process, stay still, keep their eyes open, and breathe normally during the acquisition period in order to acquire HRV data.

2.4. Revised Competitive State Anxiety Inventory 2 (CSAI-2R)

To evaluate pre-competitive anxiety, participants completed the Revised Competitive State Anxiety Inventory 2 (CSAI-2R) questionnaire. This instrument, developed by Cox *et al.* [40], is a condensed version of the CSAI-2 proposed by Martens *et al.* [41]. The CSAI-2R comprises 17 questions categorized into three components: cognitive anxiety (7 items), somatic anxiety (5 items), and self-confidence (5 items). The cumulative score for each sub-scale is determined by summing the Likert points obtained for each item. Each question offers



Fig. (1). Experimental design.

four response alternatives using a Likert scale ranging from 1 (strongly disagree) to 4 (strongly agree). The Brazilian adaptation of CSAI-2R demonstrated favorable psychometric properties for evaluating pre-competitive anxiety in Brazilian athletes [39]. The assessment duration was limited to 10 minutes. In our sample, the Cronbach's α values were 0.76, 0.81, and 0.79 for the sub-scales of cognitive anxiety, somatic anxiety, and self-confidence, respectively.

2.5. Heart Rate Variability

HRV information of all players was recorded and analyzed while they were seated on a bench in a room with air conditioning [42]. Once the assessment of competitive anxiety was completed (approximately 5 minutes), the players were provided with moistened strap transmitters, which were securely fastened to their chests. Subsequently, the players verified that the receiver of the heart rate monitor was functioning properly in order to acquire RR intervals [43]. The HRV measurements were taken at rest using a Polar V800 cardiotachometer (Polar[™], Kempele, Finland) with a sampling rate of 1,000 Hz [44]. HRV was recorded for 10 minutes, which included 5 minutes for stabilization and five minutes for poststabilization. Using specialized software, data pertaining to the 5 minutes following stabilization was collected and downloaded for analysis (Polar Precision Performance, PolarTM, Kempele, Finland). The 5-minute poststabilization HRV indicators were examined using the KubiosTM HRV program (Biomedical Signal Analysis Group, Department of Applied Physics, University of Kuopio, Kuopio, Finland) [45]. Artifacts ($\leq 2\%$) in the data were visually inspected and manually removed with the use of interpolated adjacent RR interval values (filter power < medium) [46]. The variables were reliant on the time domain (standard deviation of the mean of the qualifying NN interval) and frequency domain (high frequency (HF) and sympatho-vagal balance (LF/HF)).

2.6. Statistical Analyses

Shapiro-Wilk and Levene's tests were carried out to check the homogeneity and normality, respectively. As observed in the normality tests, the data followed a normal distribution. The information was displayed using mean and standard deviation ($M \pm SD$). At the initial stage (BL), the criteria for age, weight, height, time of experience, CSAI-2R, and HRV were satisfied. Consequently, independent samples t-tests were carried out to compare distinctions between the two groups (VG vs DG) at the BL. A 2 \times 2 mixed-factor analysis of variance (ANOVA) was utilized to determine variances between the VG and DG (between-group effects) and changes across the baseline. pre-game, and post-game periods (within-group effects) for CSAI-2R and HRV measurements in both the time and frequency domains. Subsequent analysis was performed using the Bonferroni method to assess the impacts within each group. The significance level was established at 5% (p < 0.05). Effect sizes were computed and interpreted according to Cohen's d guidelines: 0.00 to 0.19 (trivial), 0 .20 to 0 .49 (small), 0 .50 to 0 .79 (moderate), and ≥ 0.80 (large) [47]. Pearson's correlation analysis (bivariate) was carried out to examine the relationship between CSAI-2R and HRV measures [48]. All statistical analyses were performed using GraphPad Prism software, version 8.0.1.

3. RESULTS

The analyzed groups exhibited homogeneity as there were no significant differences between the groups in terms of age ($25.04\pm2.77 vs. 24.92\pm2.41$ years old, p = 0.87), weight ($79.6 \pm 2.1 vs. 77.4 \pm 2.6$ Kg, p = 0.78), height ($179.5 \pm 1.6 vs. 177.7 \pm 1.3$ cm, p = 0.69), time of experience ($7.92 \pm 1.29 vs. 7.44\pm1.38$ years, p = 0.22), and physical activity level ($37 \pm 1.2 vs. 35 \pm 0.9$ min per week, p = 0.63).

3.1. Revised Competitive State Anxiety Inventory 2 (CSAI-2R)

There were no significant differences between groups in terms of self-confidence at the baseline (p = 0.999) and pregame (p = 0.184) periods. Additionally, there were no withingroup differences between the baseline and pre-game times for the VG (p = 0.754) and the DG (p = 0.844). Nevertheless, in the post-game period, the Competitive State Anxiety Inventory-2 Revised (CSAI-2R) scores were higher in the VG group compared to the DG group ($p \le 0.0001$; effect size d = 2.26, 95% confidence interval: 1.52 to 2.93, (Fig. **2A**). A mixed analysis of variance revealed a significant interaction between group and time (F(2, 144) = 370.7; $p \le 0.0001$), main effects for the group (F(1, 144) = 313.4; $p \le 0.0001$), and time (F(2, 144) = 47.28; p = 0.003) for anxiety state.



Fig. (2). Anxiety patterns in both groups.

(A) *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to post-game time ($p \le 0.0001$), #*Significant difference compared to BL and pre-game times ($p \le 0.0001$); (B) *Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$); (C) *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$), #Significant difference compared to BL and pre-game times ($p \le 0.0001$).

The interaction analysis showed increased levels of self-confidence post-game (17.40 \pm 1.15) compared to BL (9.79 \pm 1.53) and pre-game (10.78 \pm 0.92) in the VG (p \leq 0.0001; d = 5.62, CI 95%: 4.32 to 6.75 and d = 6.36, CI 95%: 4.91 to 7.60, respectively). On the other hand, there was a decreased self-confidence score post-game (8.1 \pm 5.7) compared to BL (10.52 \pm 1.12) and pre-game (10.79 \pm 1.53) in the DG (p \leq 0.0001; d = 0.59, CI 95%: 0.01 to 1.15 and d = 0.64, CI 95%: 0.07 to 1.20, respectively).

For somatic anxiety, no differences were found between groups at BL (p = 0.827) and pre-game (p =0.265). Additionally, no intra-group differences were found between BL and pre-game in the VG (p = 0.867) and DG (p= 0.215). CSAI-2R in the post-game time was lower in VG than in DG ($p \le 0.0001$; d = 9.59, CI 95%: 7.51 to 11.37, Fig. 2B). The mixed analysis of variance showed a significant group-by-time interaction (F(2, 144) = 309.4; p \leq 0.0001), main effects for group (F(1, 144) = 387.9; p \leq 0.0001), and time (F(2, 144) = 22.08; $p \le 0.0001$) for anxiety state. The interaction analysis revealed a reduction in somatic anxiety levels post-game (6.16 \pm 0 .86) compared to BL (9.58 \pm 0.88) and pre-game (10.04 \pm 0.90) in the VG ($p \le 0.0001$; d = 3.93, CI 95%: 2.93 to 4.81 and d = 4.41, CI 95%: 3 .32 to 5. 35, respectively). In contrast, there was an increased somatic anxiety score post-game (16.12 \pm 1 .19) compared to BL (9.79 \pm 1 .10) and pre-game (10.54 \pm 1.58) in the DG (p \leq 0.0001; d = 5.52, CI 95%: 4.24 to 6.64 and d = 3 .99, CI 95%: 2.98 to 4.88, respectively).

For cognitive anxiety, no differences were found between groups at BL (p = 0.999) and pre-game (p =0.999) times. Additionally, no intra-group differences were found between BL and pre-game times in the VG (p =(0.492) and DG (p = (0.815)). CSAI-2R in the post-game time was lower in VG than in DG ($p \le 0.0001$; d = 8.42, CI 95%: 6.57 to 10.00, Fig. 2C). The mixed analysis of variance revealed a significant group-by-time interaction (F (2, 144) = 237.6; $p \le 0.0001$), main effects for group (F(1, 144) = 265.4; $p \le 0.0001$), and time (F(2, 144) = 53.62; $p \le$ 0.0001) for anxiety state. The interaction analysis showed a reduction in cognitive anxiety levels post-game (6.70 \pm 1.19) compared to BL (9.16 \pm 1.12) and pre-game (9.95 \pm 1.78) in the VG ($p \le 0.0001$; d = 2.13, CI 95%: 1.41 to 2.79 and d = 2.15, CI 95%: 1.42 to 2.80, respectively). Conversely, there was an increased cognitive anxiety score post-game (17.37 \pm 1.34) compared to BL (9.37 \pm 1.05) and pre-game (10.04 \pm 1.75) in the DG (p \leq 0.0001; d = 6.65, CI 95%: 5.14 to 7.94 and d = 4.70, CI 95%: 3.57 to 5.69, respectively).

3.2. Heart Rate Variability (HRV)

In the time domain, at baseline and pre-game, there were no differences in SDNN (standard deviation of normal-to-normal intervals) between the groups. However, SDNN was higher in the VG than the DG after the game ($p \le 0.001$; d = 3.73, CI 95%: 2.76 to 4.58). There was a significant group-by-time interaction (F (2, 144) = 39,82; $p \le 0.001$), with SDNN increasing in the VG ($p \le 0.001$; d = 2.13, CI 95%: 1.41 to 2.79 and d = 2.06, CI 95%: 1.35 to

2.71, respectively) but decreasing in the DG (p = 0.001; d = 1.05, CI 95%: 0.44 to 1.62 and p \leq 0.001; d = 1.22, CI 95%: 0.61 to 1.81, respectively) from baseline and pregame to post-game (Fig. **3A**).

Similar patterns were observed for rMSSD, which was higher in VG than in DG ($p \le 0.001$; d = 2.79, CI 95%: 1.97 to 3.52). There was a significant group-by-time interaction (F (2, 144) = 50,92; $p \le 0.001$), with rMSSD increasing in the VG ($p \le 0.001$; d = 1.27, CI 95%: 0.65 to 1.86 and p = 0.003; d = 2.15, CI 95%: 1.43 to 2.81, respectively) but decreasing in the DG ($p \le 0.001$; d = 1.82, CI 95%: 1.14 to 2.45 and d = 1.74, CI 95%: 1.06 to 2.36, respectively) from baseline and pre-game to postgame (Fig. **3B**).

Considering frequency domain measures, at baseline and pre-game, there were no differences in HF (high frequency) power between the groups. HF power was higher in the VG than the DG after the game ($p \le 0.0001$; d = 5.09, CI 95%: 3.89 to 6.14). There was a significant group-by-time interaction (*F* (2, 144) = 72,24; $p \le 0.001$), with HF increasing in the VG ($p \le 0.001$; d = 1.57, CI 95%: 0.92 to 2.18 and d = 2.20, CI 95%: 1.47 to 2.87, respectively) but decreasing in the DG ($p \le 0.001$; d = 2.33, CI 95%: 1.58 to 3.01 and d = 3.09, CI 95%: 2.23 to 3.86, respectively) from baseline/pre-game to post-game (Fig. 3C). For LF/HF (low frequency/high frequency) ratio, there were no differences between groups at baseline and pre-game, but it was lower in the VG than the DG after the game ($p \le 0.001$; d = 2.59, CI 95%: 1.80 to 3.30). There was a significant group-by-time interaction (F(2, 144) =27,55; $p \le 0.001$), with decreasing in LF-HF in the VG ($p \le$ 0.001; d = 1.93, CI 95%: 1.23 to 2.57 and d = 1.98, CI 95%: 1.28 to 2.62, respectively), but increasing in the DG (p = 0.004; d = 0.95, CI 95%; 0.35 to 1.52 and p = 0.002;d = 1.03, CI 95%: 0.43 to 1.61, respectively) from baseline/pre-game to post-game (Fig. 3D).

The study did not find any significant correlations between the HRV measures and the CSAI-2R (competitive state anxiety inventory) scores.

4. DISCUSSION

This current study aimed to investigate competitive anxiety and HRV responses before and after successful and unsuccessful matches (i.e., victory and defeat) in professional esports players. According to our hypothesis, winners had lower cognitive and somatic anxiety, higher self-confidence scores, and better HRV responses in comparison to losers who had the opposite results after the match.

Several studies have discussed the importance of monitoring anxiety and HRV in athletes. However, most studies focused on the impact of results in pre-competitive anxiety [23-26, 28, 49], HRV [27], or competitive anxiety responses before and after a competition [28], but not on the effects of results on competitive anxiety and HRV responses jointly. In line with that, our study revealed that victory and defeat interfere with HRV and anxiety responses.



Fig. (3). HRV in both groups. (**A**) SDNN: *Significant difference compared to BL and pre-game times ($p \le 0.0001$). *Significant difference compared to post-game time ($p \le 0.0001$), *Significant difference compared to BL and pre-game times (p = 0.001 and $p \le 0.0001$, respectively); (**B**) rMSSD: *Significant difference compared to BL and pre-game times ($p \le 0.0001$; p = 0.003, respectively), *Significant difference compared to BL and pre-game times ($p \le 0.0001$); (**C**) HF: *Significant difference compared to BL and pre-game times ($p \le 0.0001$); (**C**) HF: *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to post-game time ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$); (**D**) LF-HF: *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$); *Significant difference compared to BL and pre-game times ($p \le 0.0001$); *Significant difference compared to BL and pre-game times ($p \le 0.0001$); *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times ($p \le 0.0001$), *Significant difference compared to BL and pre-game times (p

Regarding anxiety responses, our findings indicated that VG experienced significantly reduced cognitive and somatic anxiety scores, along with increased selfconfidence scores at the post-game time compared to baseline (BL) and pre-game times. In contrast, DG showed the opposite pattern, with significantly increased cognitive and somatic anxiety scores, as well as significantly reduced self-confidence scores at the post-game time compared to BL and pre-game times. Furthermore, there were no differences between BL and pre-game times, as well as between BL and post-game in both VG and DG on the Competitive State Anxiety Inventory 2 Revised (CSAI-2R).

Concerning heart rate variability (HRV), our results revealed a substantial increase in RR, SDNN, rMSSD, pNN50, and HF and a substantial decrease in LF and LF/HF in the VG. DG demonstrated a substantial decrease in RR, SDNN, rMSSD, and HF, and a substantial increase in LF and LF/HF. Moreover, there were no differences in BL and pre-game times, as well as between BL and postgame times in both VG and DG in any HRV measure. Our results align with the existing literature on HRV and competitive anxiety, emphasizing that different magnitudes of response may occur depending on the situation because of the robust correlation between the autonomic nervous system (ANS) and anxiety.

The results of a competition are considered a significant factor contributing to anxiety responses and changes. Additionally, recent studies have HRV underscored the importance of using HRV to monitor players, and the assessment of anxiety levels in highperformance sports is deemed crucial for emotional control throughout the competition. To the best of our knowledge, this study is the first to investigate the impact of victory and defeat on heart rate variability (HRV) responses and competitive anxiety. Despite this, it is reasonable to suggest that victory and defeat have positive and negative effects on HRV and anxiety behaviors. In this context, our results regarding anxiety levels are in line with a previous study by Fuentes-Garcia et al. [28], even though they specifically focused on competitive anxiety. The authors explored the influence of victory and defeat on pre-competitive anxiety in athletes during international tennis competitions. Significant differences were noted in

cognitive anxiety and self-assurance levels when analyzing pre-game and post-game results among winners and defeated players. Winners encountered a notable decrease in cognitive anxiety and a rise in self-confidence following the match.

These findings reinforce our results, suggesting that esports players who emerge victorious tend to exhibit more adaptive responses in cognitive and somatic anxiety, self-confidence, and greater parasympathetic activation compared to losers. These differences can extend to HRV. In a similar vein, a study by Pena [27] demonstrated that junior tennis players who were defeated experienced a significant reduction in PNN50 after the match compared to the winners. These findings suggest that losers had higher sympathetic activity than winners after the match. Overall, anxiety can be regarded as a failure in the ability to inhibit cognitive, affective, behavioral, and physiological responses, leading to reduced vagal outflow and lowered HRV.

The neurovisceral integration model emphasizes the involvement of the prefrontal cortex in inhibitory processes mediated by the vagus nerve, which can be observed through measures of heart rate variability (HRV) [16]. An imbalance between sympathetic and parasympathetic activity, which is linked to task performance [50, 51], can give rise to significant alterations in prefrontal cortex functioning (responsible for executive functions) at various levels [52]. Consequently, these changes can impact an individual's psychological state, including the experience of anxiety [53].

There is a correlation between cognitive functioning, anxiety levels, and heart rate variability (HRV), as the connection between the brain and heart is constant and crucial for efficient and regulated functioning [54]. However, anxiety levels impact HRV through the vagus nerve (parasympathetic), which influences the sinoatrial node of the heart, resulting in an imbalance between sympathetic and parasympathetic activity [53, 55]. Moreover, given that players are consistently active, whether during training or competition, their performance can influence neurocardiac dynamics, and the outcome of a match (be it a win or a loss) can impact the establishment of a sympathetic/parasympathetic balance [50]. In esports, sustained demands on concentration, attention, and decision-making are required throughout the match to ensure task efficiency [8], and this demand leads to changes in anxiety levels and HRV, influencing positivelv or negatively on match performance. Furthermore, the concept of the polyvagal theory [56] emphasizes the significance of the myelinated vagus nerve in facilitating social interaction and communication, as well as inducing states of relaxation by suppressing sympathetic activity in the heart and modulating the hypopituitary-adrenal axis. When an individual perceives their environment as safe, there is an increase in vagal outflow, which promotes restoration, maintenance of internal balance, and engagement in social interactions. However, impairments in these neural processes can lead to possible damage to the perception that certain environments are safe, as well as difficulties in communication and social inter-action, which can impair the performance of players during training and matches. Therefore, the link between anxiety and deficits in the inhibition of sympathetic function, as well as the difficulty in communication and social interaction, can be explained by the polyvagal theory.

Our findings have important implications for establishing the relationship between the reduction in HRV parameters with anxiety in professional esports players. Based on clinical studies, individuals with anxiety disorders have a higher attentional bias related to threatening cues when compared to individuals without anxiety disorders [57]. This pattern of behavior results in consistently elevated levels of corticotropin (released in response to stress) and cortisol, leading to a persistent suppression of parasympathetic activity (i.e., low vagal tone) [58]. Moreover, ongoing concerns may disrupt the regulation of cardiac autonomic function in the presence of threatening stimuli [14]. It is plausible that a decline in vagal function (which plays a critical role in regulating the hypothalamic-pituitary-adrenal axis) [59] contributes to heightened reactivity to stressors.

The inability to disengage from threat detection intensifies sympathetic activation due to the enduring reduction of parasympathetic activity (and subsequent long-term decreases in heart rate variability). Machado et al. [7] explored for the first time the impact of victory and defeat in playoff matches on competitive anxiety and HRV in professional esports players. Hence, more studies are required to better comprehend HRV patterns and emotional states (beyond stress and anxiety) in professional esports players, both prior to and following different game outcomes, such as victories, defeats, and potential draws. These studies hold significance in comprehending the dynamics of these factors over the course of a competitive season, thereby offering valuable insights for potential interventions aimed at facilitating the players' recovery between matches and throughout the entire competition.

The present study uncovered noteworthy, albeit preliminary, discoveries that hold relevance for professional esports players and coaches. Our results indicated that victory and defeat in playoffs provoke different responses in HRV and anxiety state. As a practical implication, we can use HRV as a monitoring tool for athletes during competition, especially in playoffs. The use of HRV in quantifying training loads is already established [31], as well as the analysis and organization of training programs [34]. This enables the evaluation, monitoring, recovery, and improvement of performance [60]. However, HRV was found to respond in accordance with the level of competitive anxiety influenced by environmental factors, such as victory or defeat. Thus, HRV plays an important role in this type of monitoring, as it is influenced by the emotional state [59].

In line with this, rMSSD is the most used HRV indicator at rest [61], as it is effective in identifying the parasympathetic level of the heart. Therefore, esports

athletes and coaches should consider these results, as they can be utilized to evaluate, track, restore, and uphold effectiveness in training and game execution. This study has certain limitations. The examination of competitive anxiety and heart rate variability (HRV) in esports players during the game was not carried out, just before and after the games. Intra-match HRV and anxiety can provide insights into the athlete's psychophysiological state during the match. Without this data, the study may overlook critical fluctuations in stress and recovery that occur in real time, potentially skewing the relationship between HRV and performance outcomes. Moreover, HRV and anxiety are often used as a marker of fatigue and recovery. It is difficult to assess how fatigue accumulates throughout the match and how it correlates with performance metrics, leading to an incomplete picture of an athlete's condition. Also, there was no comparison of competitive anxiety and HRV parameters among players with different functions in the team. Furthermore, there was no assessment of a protective factor through psychotherapy or counseling that could prevent or alleviate symptoms early on.

For future studies, we recommend investigating the effects of ergogenic aids to enhance both neurocognitive and neuromotor performance, utilizing substances, such as caffeine [62] and non-invasive brain stimulation techniques like transcranial direct current stimulation (tDCS) [8, 63]. Furthermore, it would be beneficial to consider future studies that include measures during the matches and analyze differences according to team roles.

CONCLUSION

This study aimed to analyze the behavior of competitive anxiety and HRV in different teams during playoff matches in professional esports players. Our results revealed that VG exhibited better HRV responses, indicating greater parasympathetic activation. VG also showed lower levels of cognitive and somatic anxiety and higher levels of self-confidence in the post-game time. In contrast, DG demonstrated worse HRV responses, indicating greater sympathetic activation, along with higher levels of cognitive and somatic anxiety and lower levels of self-confidence in the same post-game period. Future research should delve into analyzing HRV behavior and other emotional variables before, during, and after the conclusion of competitive esports matches with different outcomes (victory, defeat, and perhaps draw). Additionally, exploring the impact of various ergogenic aids on neurocognitive and neuromotor performance would be valuable.

AUTHORS' CONTRIBUTIONS

S.M.: Writing the Paper; L.D.O.S.A., L.C., A.E.N.: Data Analysis or Interpretation; D.S.T., B.T., D.M.: Study Concept or Design.

LIST OF ABBREVIATIONS

- HRV = Heart Rate Variability
- VG = Victory Group

DG = Defeated Group

CSAI-2R = Revised Competitive State Anxiety Inventory 2

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

This study was approved by the Ethics Committee of the University of Beira Interior, Portugal (approval number: D2574).

HUMAN AND ANIMAL RIGHTS

All procedures performed in studies involving human participants were in accordance with the ethical standards of institutional and/or research committee and with the 1975 Declaration of Helsinki, as revised in 2013.

CONSENT FOR PUBLICATION

The participants provided their written informed consent to participate in this study.

AVAILABILITY OF DATA AND MATERIALS

The data sets used and/or analysed during this study are available from the corresponding author [S.M.N] upon request.

FUNDING

This research was supported by national funds through the Portuguese Foundation for Science and Technology, I.P., Portugal under the project UID04045/2020.

CONFLICT OF INTEREST

Drs. Antonio Nardi and Sergio Machado are editorial advisory board members of the journal Clinical Practice & Epidemiology in Mental Health.

ACKNOWLEDGEMENTS

Declared none.

8542197

REFERENCES

- [1] Paridon VKN, Timmis MA, Nevison CM, Bristow M. The anticipatory stress response to sport competition; a systematic review with meta-analysis of cortisol reactivity. BMJ Open Sport Exerc Med 2017; 3(1): e000261.
 - http://dx.doi.org/10.1136/bmjsem-2017-000261 PMID: 29177073
- [2] Ford J, Ildefonso K, Jones M, Arvinen-Barrow M. Sport-related anxiety: Current insights. Open Access J Sports Med 2017; 8: 205-12.
 - http://dx.doi.org/10.2147/OAJSM.S125845 PMID: 29138604
- [3] Jones G. More than just a game: Research developments and issues in competitive anxiety in sport. Br J Psychol 1995; 86(4): 449-78. http://dx.doi.org/10.1111/j.2044-8295.1995.tb02565.x
- [4] Hanton S, Neil R, Mellalieu SD. Recent developments in competitive anxiety direction and competition stress research. Int Rev Sport Exerc Psychol 2008; 1(1): 45-57. http://dx.doi.org/10.1080/17509840701827445
- [5] Jordet G, Gemser ME, Lemmink KP. Perceived control and anxiety. Int J Sport Psychol 2006; 37: 281-98.
- [6] López-Torres M, Torregrosa M, Roca J. Characteristics of "flow", anxiety and emotional state, in relation to the performance of elite athletes. Cuad Psicol Deporte 2007; 7: 25-44.

- [7] Machado S, Sant'Ana LO, Cid L, et al. Impact of victory and defeat on the perceived stress and autonomic regulation of professional eSports athletes. Front Physiol 2022; •••: 13.
- [8] Machado S, Travassos B, Teixeira DS, Rodrigues F, Cid L, Monteiro D. Could tDCS Be a potential performance-enhancing tool for acute neurocognitive modulation in eSports? A perspective review. Int J Environ Res Public Health 2021; 18(7): 3678.

http://dx.doi.org/10.3390/ijerph18073678 PMID: 33916018

[9] Freeman R. Assessment of cardiovascular autonomic function. Clin Neurophysiol 2006; 117(4): 716-30.

http://dx.doi.org/10.1016/j.clinph.2005.09.027 PMID: 16464634

- [10] Paniccia M, Paniccia D, Thomas S, Taha T, Reed N. Clinical and non-clinical depression and anxiety in young people: A scoping review on heart rate variability. Auton Neurosci 2017; 208: 1-14. http://dx.doi.org/10.1016/j.autneu.2017.08.008 PMID: 28870754
- [11] Blásquez CJC, Font RG, Ortís CL. Heart-rate variability and precompetitive anxiety in swimmers. Psicothema 2009; 21(4): 531-6.

PMID: 19861094

- [12] Ayuso-Moreno R, Fuentes-García JP, Collado-Mateo D, Villafaina S. Heart rate variability and pre-competitive anxiety according to the demanding level of the match in female soccer athletes. Physiol Behav 2020; 222: 112926. http://dx.doi.org/10.1016/j.physbeh.2020.112926 PMID: 32407830
- [13] Mateo M, Blasco-Lafarga C, Martínez-Navarro I, Guzmán JF, Zabala M. Heart rate variability and pre-competitive anxiety in BMX discipline. Eur J Appl Physiol 2012; 112(1): 113-23. http://dx.doi.org/10.1007/s00421-011-1962-8 PMID: 21503698
- [14] Berntson GG, Sarter M, Cacioppo JT. Anxiety and cardiovascular reactivity: The basal forebrain cholinergic link. Behav Brain Res 1998; 94(2): 225-48.

http://dx.doi.org/10.1016/S0166-4328(98)00041-2 PMID: 9722275

- [15] Berntson GS, Cacioppo JT. Psychophysiology.Biological Psychiatry. Hoboken, New Jersey: John Wiley & Sons 2002. http://dx.doi.org/10.1002/0470854871.chix
- [16] Friedman BH. An autonomic flexibility-neurovisceral integration model of anxiety and cardiac vagal tone. Biol Psychol 2007; 74(2): 185-99.

http://dx.doi.org/10.1016/j.biopsycho.2005.08.009 PMID: 17069959

- [17] Grossman P, Taylor EW. Toward understanding respiratory sinus arrhythmia: Relations to cardiac vagal tone, evolution and biobehavioral functions. Biol Psychol 2007; 74(2): 263-85. http://dx.doi.org/10.1016/j.biopsycho.2005.11.014 PMID: 17081672
- [18] Porges SW. A phylogenetic journey through the vague and ambiguous Xth cranial nerve: A commentary on contemporary heart rate variability research. Biol Psychol 2007; 74(2): 301-7. http://dx.doi.org/10.1016/j.biopsycho.2006.08.007 PMID: 17055142
- [19] Dishman RK, Nakamura Y, Garcia ME, Thompson RW, Dunn AL, Blair SN. Heart rate variability, trait anxiety, and perceived stress among physically fit men and women. Int J Psychophysiol 2000; 37(2): 121-33. http://dx.doi.org/10.1016/S0167-8760(00)00085-4 PMID:

10831999

[20] Hjortskov N, Rissén D, Blangsted AK, Fallentin N, Lundberg U, Søgaard K. The effect of mental stress on heart rate variability and blood pressure during computer work. Eur J Appl Physiol 2004; 92(1-2): 84-9. http://dx.doi.org/10.1007/s00421-004-1055-z PMID: 14991326

http://dx.doi.org/10.100//s00421-004-1055-2 PMID: 14991520

- [21] Ruediger H, Seibt R, Scheuch K, Krause M, Alam S. Sympathetic and parasympathetic activation in heart rate variability in male hypertensive patients under mental stress. J Hum Hypertens 2004; 18(5): 307-15. http://dx.doi.org/10.1038/sj.jhh.1001671 PMID: 15103310
- [22] Cohen H, Benjamin J. Power spectrum analysis and cardiovascular morbidity in anxiety disorders. Auton Neurosci 2006; 128(1-2): 1-8.

http://dx.doi.org/10.1016/j.autneu.2005.06.007 PMID: 16731048

- [23] Covassin T, Pero S. The relationship between self-confidence, mood state, and anxiety among collegiate tennis players. J Sport Behav 2004; 27: 230-42.
- [24] Fernandez-Fernandez J, Boullosa DA, Sanz-Rivas D, Abreu L, Filaire E, Mendez-Villanueva A. Psychophysiological stress responses during training and competition in young female competitive tennis players. Int J Sports Med 2015; 36(1): 22-8. PMID: 25251448
- [25] Filaire E, Alix D, Ferrand C, Verger M. Psychophysiological stress in tennis players during the first single match of a tournament. Psychoneuroendocrinology 2009; 34(1): 150-7. http://dx.doi.org/10.1016/j.psyneuen.2008.08.022 PMID: 18845399
- [26] Fratta LI, Franceschelli S, Speranza L, et al. Salivary oxytocin, cognitive anxiety and self-confidence in pre-competition athletes. Sci Rep 2021; 11(1): 16877. http://dx.doi.org/10.1038/s41598-021-96392-7 PMID: 34413428
- [27] Pena J. Heart rate variability as a marker of psychobiological stress and anxiety in elite junior tennis players in Venezuela. J Am Coll Cardiol 2020; 75(11): 1613. http://dx.doi.org/10.1016/S0735-1097(20)32240-3
- [28] Fuentes-García JP, Villafaina S, Martínez-Gallego R, Crespo M. Crespo M. Pre- and post-competitive anxiety and match outcome in elite international junior tennis players. Int J Sport Sci Coach 2022.http://journals.sagepub.com/doi/10.1177/174795412211223 96
- [29] Pereira MA, Costa JA, Verhagen E, Figueiredo P, Brito J. Associations between sports participation and health: A Scoping Review. Sports Med 2022; 52(9): 2039-60. http://dx.doi.org/10.1007/s40279-022-01684-1 PMID: 35486374
- [30] Trotter MG, Coulter TJ, Davis PA, Poulus DR, Polman R. Social support, self-regulation, and psychological skill use in E-athletes. Front Psychol 2021; 12: 722030. http://dx.doi.org/10.3389/fpsyg.2021.722030 PMID: 34858261
- [31] Borresen J, Lambert IM. The quantification of training load, the training response and the effect on performance. Sports Med 2009; 39(9): 779-95. http://dx.doi.org/10.2165/11317780-000000000-00000 PMID: 19691366

[32] Morgan JS, Mora MJA. Effect of heart rate Variability biofeedback on sport performance: A Systematic review. Appl Psychophysiol Biofeedback 2017; 42(3): 235-45.

http://dx.doi.org/10.1007/s10484-017-9364-2 PMID: 28573597 [33] Pusenjak N, Grad A, Tusak M, Leskovsek M, Schwarzlin R. Can

biofeedback training of psychophysiological responses enhance athletes' sport performance? A practitioner's perspective. Phys Sportsmed 2015; 43(3): 287-99. http://dx.doi.org/10.1080/00913847.2015.1069169 PMID:

26200172

- [34] Tønnessen E, Sylta Ø, Haugen TA, Hem E, Svendsen IS, Seiler S. The road to gold: Training and peaking characteristics in the year prior to a gold medal endurance performance. PLoS One 2014; 9: e101796.
- [35] Makivić B, Nikić DM, Willis MS. Heart rate variability (HRV) as a tool for diagnostic and monitoring performance in sport and physical activities. JEPonline 2013; 16: 103131.
- [36] Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. Behav Res Methods 2009; 41(4): 1149-60. http://dx.doi.org/10.3758/BRM.41.4.1149 PMID: 19897823
- [37] Oliveira-Silva I, Silva VA, Cunha RM, Foster C. Autonomic changes induced by pre-competitive stress in cyclists in relation to physical fitness and anxiety.PLoS One 2018; 13(12): e0209834. http://dx.doi.org/10.1371/journal.pone.0209834
- [38] Hallal PC, Victora CG. Reliability and validity of the international physical activity questionnaire (IPAQ). Med Sci Sports Exerc 2004; 36(3): 556. http://dx.doi.org/10.1249/01.MSS.0000117161.66394.07 PMID: 15076800

- [39] Fernandes MG, Nunes SA, Raposo JV, Fernandes HM, Brustad R. The CSAI-2: An examination of the instrument's factorial validity and reliability of the intensity, direction and frequency dimensions with brazilian athletes. J Appl Sport Psychol 2013; 25(4): 377-91. http://dx.doi.org/10.1080/10413200.2012.744780
- [40] Cox RH, Martens MP, Russell WD. Measuring anxiety in athletics: The revised competitive state anxiety inventory - 2. J Sport Exerc Psychol 2003; 25(4): 519-33. http://dx.doi.org/10.1123/jsep.25.4.519
- [41] Martens R, Burton D, Vealey RS, Bump LA, Smith DE. Development and validation of the competitive state anxiety inventory-2 (CSAI-2). Competitive Anxiety in sport. Chapaign: Human Kinetics 1990; pp. 117-90.
- [42] Holmes CJ, Fedewa MV, Dobbs WC, et al. The effects of different body positions on the accuracy of ultra-short-term heart rate variability indexes. J High Technol Manage Res 2020; 31(1): 100375.

http://dx.doi.org/10.1016/j.hitech.2020.100375

- [43] Geus DEJC, Gianaros PJ, Brindle RC, Jennings JR, Berntson GG. Should heart rate variability be "corrected" for heart rate? Biological, quantitative, and interpretive considerations. Psychophysiology 2019; 56(2): e13287. http://dx.doi.org/10.1111/psyp.13287 PMID: 30357862
- [44] Quintana DS, Heathers JAJ, Kemp AH. On the validity of using the Polar RS800 heart rate monitor for heart rate variability research. Eur J Appl Physiol 2012; 112(12): 4179-80. http://dx.doi.org/10.1007/s00421-012-2453-2 PMID: 22790488
- [45] Draghici AE, Taylor JA. The physiological basis and measurement of heart rate variability in humans. J Physiol Anthropol 2016; 35(1): 22. http://dx.doi.org/10.1186/s40101-016-0113-7 PMID: 27680542
- [46] Johnston BW, Barrett-Jolley R, Krige A, Welters ID. Heart rate variability: Measurement and emerging use in critical care medicine. J Intensive Care Soc 2020; 21(2): 148-57. http://dx.doi.org/10.1177/1751143719853744 PMID: 32489411
- [47] Cohen J. Statistical power analysis for the behavioral sciences. New York: NY Routledge Acad. 1988.
- [48] Schober P, Boer C, Schwarte LA. Correlation coefficients: Appropriate use and interpretation. Anesth Analg 2018; 126(5): 1763-8. http://dx.doi.org/10.1213/ANE.000000000002864 PMID:
- 29481436 [49] Terry PC, Cox JA, Lane AM, Karageorghis CI. Measures of anxiety among tennis players in singles and doubles matches. Percept Mot Skills 1996; 83(2): 595-603.

http://dx.doi.org/10.2466/pms.1996.83.2.595

[50] Britton DM, Kavanagh EJ, Polman RCJ. Validating a self-report measure of student athletes' perceived stress reactivity: Associations with heart-rate variability and stress appraisals. Front Psychol 2019; 10: 1083. http://dx.doi.org/10.3389/fpsyg.2019.01083 PMID: 31143153

[51] Barnes CM, Dyne VL. 'I'm tired': Differential effects of physical and emotional fatigue on workload management strategies. Hum Relat 2009; 62(1): 59-92.

http://dx.doi.org/10.1177/0018726708099518

[52] Ai JY, Chen FT, Hsieh SS, et al. The effect of acute high-intensity

interval training on executive function: A systematic review. Int J Environ Res Public Health 2021; 18(7): 3593.

- http://dx.doi.org/10.3390/ijerph18073593 PMID: 33808399 [53] Kemp AH, Quintana DS. The relationship between mental and
- physical health: Insights from the study of heart rate variability. Int J Psychophysiol 2013; 89(3): 288-96. http://dx.doi.org/10.1016/j.ijpsycho.2013.06.018 PMID: 23797149
- [54] Lin F, Heffner KL, Ren P, Tadin D. A role of the parasympathetic nervous system in cognitiva training. Curr Alzheimer Res 2017; 14(7): 784-9. PMID: 28164771
- [55] Quintana DS, Guastella AJ, Outhred T, Hickie IB, Kemp AH. Heart rate variability is associated with emotion recognition: Direct evidence for a relationship between the autonomic nervous system and social cognition. Int J Psychophysiol 2012; 86(2): 168-72.

http://dx.doi.org/10.1016/j.ijpsycho.2012.08.012 PMID: 22940643

- [56] Porges SW. The Polyvagal theory: Neurophysiological foundations of emotions, attachment, communications and self-regulation. 2011. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC3490536/
- [57] Bar-Haim Y, Lamy D, Pergamin L, Bakermans-Kranenburg MJ, IJzendoorn VMH. Threat-related attentional bias in anxious and nonanxious individuals: A meta-analytic study. Psychol Bull 2007; 133(1): 1-24.

http://dx.doi.org/10.1037/0033-2909.133.1.1 PMID: 17201568

[58] Barlow DH. Unraveling the mysteries of anxiety and its disorders from the perspective of emotion theory. Am Psychol 2000; 55(11): 1247-63

http://dx.doi.org/10.1037/0003-066X.55.11.1247 PMID: 11280938

[59] Thayer JF, Sternberg E. Beyond heart rate variability: Vagal regulation of allostatic systems. Ann N Y Acad Sci 2006; 1088(1): 361-72.

http://dx.doi.org/10.1196/annals.1366.014 PMID: 17192580

[60] Meeusen R, Duclos M, Foster C, et al. Prevention, diagnosis, and treatment of the overtraining syndrome: Joint consensus statement of the European College of Sport Science and the American College of Sports Medicine. Med Sci Sports Exerc 2013; 45(1): 186-205.

http://dx.doi.org/10.1249/MSS.0b013e318279a10a PMID: 23247672

- [61] Buchheit M. Monitoring training status with HR measures: Do all roads lead to Rome? Front Physiol 2014; 5: 73. http://dx.doi.org/10.3389/fphys.2014.00073 PMID: 24578692
- [62] Sainz I, Collado-Mateo D, Coso JD. Effect of acute caffeine intake on hit accuracy and reaction time in professional e-sports players. Physiol Behav 2020; 224: 113031. http://dx.doi.org/10.1016/j.physbeh.2020.113031 PMID: 32593750
- [63] Machado S, Sant'Ana LO, Travassos B, Monteiro D. Anodal transcranial direct current stimulation reduces competitive anxiety and modulates heart rate variability in an esports player. Clin Pract Epidemiol Ment Health 2022: 18(1): e174501792209270.

http://dx.doi.org/10.2174/17450179-v18-e2209270 PMID: 37274860