

# Impact of Pre-Lockdown Hyper-Energy on Mood and Rhythm Dysregulation in Older Adults During the COVID-19 Pandemic



Diego Primavera<sup>1</sup>, Goce Kalcev<sup>1,\*</sup>, Fabrizio Bert<sup>2,3</sup>, Elisa Cantone<sup>1</sup>, Alessandra Perra<sup>1</sup>, Massimo Tusconi<sup>4</sup>, Samantha Pinna<sup>1</sup>, Germano Orrù<sup>5</sup>, Alessandra Scano<sup>5</sup>, Enzo Tramontano<sup>6</sup>, Ivan Barbov<sup>7</sup>, Marcello Nonnis<sup>8</sup>, Antonio Egidio Nardi<sup>9</sup>, Giulia Cossu<sup>1</sup>, Federica Sancassiani<sup>1</sup> and Mauro Giovanni Carta<sup>1</sup>

<sup>1</sup>Department of Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy

<sup>2</sup>Department of Public Health Sciences, University of Turin, Turin, Italy

<sup>3</sup>Collegium Medicum, University of Social Sciences, Lodz, Poland

<sup>4</sup>University Hospital of Cagliari, Cagliari, Italy

<sup>5</sup>Department of Surgical Sciences, University of Cagliari, Cagliari, Italy

<sup>6</sup>Department of Life and Environmental Sciences, University of Cagliari, Cagliari, Italy

<sup>7</sup>University Clinic of Neurology, Skopje, 1000, Skopje, North Macedonia

<sup>8</sup>Department of Pedagogy, Psychology, Philosophy, University of Cagliari, Cagliari, Italy

<sup>9</sup>Institute of Psychiatry (Ipub), Federal University of Rio De Janeiro (Ufrj), Brazil

## Abstract:

**Objective:** The aim of this work is to verify whether a cohort of elderly people with hyper-energy tended to increase depressive symptoms and misaligned social and personal rhythms during the lockdown compared to a cohort of older adults without hyper-energy one year before the lockdown.

**Methods:** The two cohorts were evaluated in April 2019 (T0) and in April 2020 (T1). Hyper-energy, cognitive performance, depressive symptoms, and social and personal rhythms were evaluated at T0 and T1.

**Results:** In the measure of the Brief Social Rhythm Scale (BSRS) score, the differences between groups in the two observation times reach statistical significance. The sub-group with previous hyper-energy at T0 but no longer having hyper-energy at T1 increases the score by more than 5 points (a higher score indicates greater rhythm dysregulation, thus having a worse regulation of rhythms at T1), while in those individuals who didn't have hyper-energy, the score remains substantially unchanged (+0.06). However, if the increase in the score from T0 and T1 is measured, both groups with hyper-energy at T0 presented a greater mean increase compared to people who did not have hyper-energy at T1. In the total of the eleven elderly people with hyper-energy at T0, the mean increase in BSRS score was  $1.05 \pm 1.19$  versus  $0.06 \pm 0.98$  ( $F=9.407$ ,  $P=0.003$ ), and in people who no longer had hyper-energy at T1, it was  $1.05 \pm 1.19$  versus  $5.50 \pm 3.83$  ( $F=105.0$ ,  $P<0.0001$ ). In people with hyper-energy at T0, the mean increase in the Patient Health Questionnaire-9 (PHQ-9) score was  $0.72 \pm 0.75$  versus  $0.01 \pm 0.28$  ( $F=37.153$ ,  $P<0.0001$ ). The gain was even higher in people who no longer had hyper-energy at T1,  $1.38 \pm 1.03$  vs.  $0.01 \pm 0.28$  ( $F=87.386$ ,  $P<0.0001$ ). An inverse linear correlation was found between energy perception (measured as the score of Item 10 of SF-12) and the score of PHQ-9 measuring depressive symptoms both at T0 and, more strongly, at T1, as well as with the BSRS scores, but only at T1.

**Conclusion:** The study, despite the limitations of a small sample, seems to confirm a greater vulnerability to the lockdown situation in people with hyperactivity, even in the absence of psychopathology (i.e., part of the bipolar spectrum).

**Keywords:** Hyper-energy, Rhythm dysregulation, advanced technologies laboratory, Depressive symptoms, COVID-19, Bipolar spectrum, Stress.

© 2025 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Received: July 13, 2024  
Revised: November 07, 2024  
Accepted: November 19, 2024  
Published: February 12, 2025

\*Address correspondence to this author at the Department of Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy; E-mail: [gocekalcev@yahoo.com](mailto:gocekalcev@yahoo.com)

*Cite as:* Primavera D, Kalcev G, Bert F, Cantone E, Perra A, Tusconi M, Pinna S, Orrù G, Scano A, Tramontano E, Barbov I, Nonnis M, Nardi A, Cossu G, Sancassiani F, Carta M. Impact of Pre-Lockdown Hyper-Energy on Mood and Rhythm Dysregulation in Older Adults During the COVID-19 Pandemic. *Clin Pract Epidemiol Ment Health*, 2025; 21: e17450179344148. <http://dx.doi.org/10.2174/0117450179344148250206065104>



Send Orders for Reprints to  
[reprints@benthamscience.net](mailto:reprints@benthamscience.net)

## 1. INTRODUCTION

The COVID-19 pandemic lockdown has placed individuals in a sort of exceptional environmental assessment, altering several risk factors for mood disorders and, among these, and not least for relevance, social and individual rhythms [1-3]. The dysregulation of individual and personal rhythms (in close connection with biorhythms) has been seen to be an element associated with mood disorders, in particular with the spectrum of bipolar disorders [4, 5]. In fact, according to some researchers, the alterations in biorhythms are considered to have a role in the genesis of mood disorders [6-8]. Also, there is evidence of a vulnerability in bipolar disorders toward triggers capable of altering social rhythms and biorhythms, such as light stimuli and light pollution [9, 10]. The circadian rhythmicity is under the control of the biological clock structures that manage the functioning of human beings under 24-hour cycles. Internal and external signals influence this clock, but light is the most important one. Socio-economic and cultural factors in modern life have led to a revolution in light patterns, which may be a cause for the increase in bipolar disorders in our society or, at least, a paradoxical stability for a strongly disadvantaging disease, and it may determine an adaptive condition of certain "subthreshold" forms [4]. A recent case-control study found that older adults with a depressive episode during the lockdown but not depressed a year earlier experienced a paradoxical increase in energy in the same year before both the lockdown and the depressive episode [11].

It was also found that non-pathological personality characteristics and traits of hyper-energy / hyperactivity (without any diagnoses of mood disorders) frequently share genetic variants that have been associated with bipolar disorder [12]. In other words, those genetic characteristics would not have been typical of bipolar disorder in "itself" but of hyper-energy, hyperactivity, and novelty-seeking traits, including people with the disorder as a sort of "tip of the iceberg" [13, 14]. Therefore, the possible onset of bipolar disorder could be understood as the convergence of those genetic variants of hyper-energy and hyperactivity with some kind of stress condition [4, 15]. It needs to be explored how factors such as different genetically specific vulnerabilities to stress, as well as the epigenetic transmission of the trauma memory and perinatal trauma, characteristics intrinsic to stress, or other environmental variables, can play a trigger for an

episode of pathological mood (in the case of the predisposition to hyper-energy, this is a path that requires entirely to be followed) [16-23]. By reversing the observation perspective of the previously cited study [11], it could be interesting to observe, according to a prospective model, how elderly people with previously assessed hyper-energy/hyperactivity were coping with the lockdown (based on their mood, personal, and social rhythms) compared to elderly people who did not show this hyper-energy or hyperactivity one year earlier, before the first peak of the pandemic.

The aim of this work is to study whether the members of a cohort of elderly people with hyperactivity/hyper-energy characteristics tended to increase depressive symptoms and regulate social and personal rhythms during the lockdown with a different profile compared to a cohort of older adults without hyperactivity/hyper-energy one year before.

A parallel evaluation was conducted to verify whether the presence of hyperactivity correlated similarly with the observed variables at the two different time points.

## 2. METHODS

### 2.1. Design and Study Sample

A cohort study was performed at the end of an RCT (Randomized-Controlled Trial) [24]. This study is a secondary analysis, meaning it utilizes data originally collected for a different primary research purpose. Specifically, the data were initially gathered for a Randomized Controlled Trial (RCT) assessing the efficacy of moderate physical exercise in elderly participants. For this study, we conducted a separate analysis on this pre-existing dataset to explore the relationship between hyper-energy and mood dysregulation during the COVID-19 lockdown. By reanalyzing the data with a different focus, we aimed to generate new insights beyond the original scope of the RCT. The study was conceived to assess, as a primary outcome, the efficacy of moderate physical exercise training. This cohort study also included the individuals selected for participation in the RCT but not involved in the final sample after the signed consensus due to over-numbering the cells of randomization. They remained on a waiting list because another trial was planned but not carried out in spite of the pandemic. As described in more detail in the previous paper [24], the criteria for inclusion included: no limitation by gender, age

≥ 65 years, living at home, and positive medical evaluation for non-competitive physical activity. On the other hand, the exclusion criteria were: age less than 65 years old, presence of health conditions causing unsuitability for moderate physical activity, living in sheltered houses or assisted residences, BMI (body mass index) > 35, history of psychosis, and presence of severe brain disease.

The sample was evaluated at the entry of RCT in April 2019 (coinciding with the entry of this cohort (T0)) and in April 2020 (end of cohort (T1) serendipity during the lockdown). Hyper-energy, cognitive performance, depressive symptoms, and social and personal rhythms were evaluated at T0 and T1.

## 2.2. Instruments

In this study, as a measure of hyper-energy, we adopted item number 10 of the Short Form Health Survey in the version of 12 items (SF-12) [25] questionnaire, “How long in the last 4 weeks did you feel full of energy?.” Only the answer number 6 on a scale from one to six (coded “always”) was taken into consideration as indicative of hyper-energy.

The self-administered tool Patient Health Questionnaire in the version with 9 items (PHQ-9) was adopted for measuring depressive symptoms [26]. The overall score of this scale is the sum of the scores of each of the items that indicate the core symptoms of the depressive episode according to the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) [27].

The functionality of social and behavioral rhythms was measured by the Italian version of the Brief Social Rhythms Scale (BSRS) [28]. A high total score on this scale indicates worse functioning.

## 2.3. Data Analysis Section

The data were analyzed using the SPSS software (version 23). The comparison between means and standard deviations of numerical data between the two cohorts at the start and end of the follow-up was conducted by means of one-way ANOVA statistics with Bonferroni corrections. The comparison at each observation point (T0 and T1) was conducted between those individuals who did not have hyper-energy at T0 (who didn't answer “6” to item 10 of the SF-12) and those who had hyper-energy at T0 (who had responded “6” to item 10 of the SF-12), and among these were also the elderly people who had hyper-energy at T0 but had no more at T1. The nominal variables were compared by the Fisher Exact test. The correlation at T0 and T1 between variables was measured by Pearson's correlation test.

In this study, hyper-energy was defined as a self-reported perception of feeling “always full of energy,” corresponding to a score of “6” on the SF-12 questionnaire item. This threshold was selected based on the study's aim to identify individuals exhibiting unusually high, sustained energy levels, a trait hypothesized to correlate with dysregulation in mood and social rhythms. While consistently high energy levels may indicate a degree of hyperactivity or heightened arousal, this classification

does not imply a diagnosis of hyperactivity disorder. Instead, it aligns with emerging evidence linking such traits to mood regulation challenges, particularly under stressors like lockdown conditions. Future studies could explore a more nuanced scale to differentiate between normative high energy and potentially maladaptive hyper-energy.

## 2.4. Ethical Aspect

The study is registered on ClinicalTrials.gov (NCT03858114). The Regional Ethical Committee has approved the study with reference number PG/2018/15546, approved 25 October 2018. This study forms part of a larger research project initially designed to evaluate the effects of moderate physical exercise on elderly participants. As part of this broader project, the data used in this secondary analysis were collected during the study period and included information relevant to physical and mental health. Although the current analysis was not initially outlined in the primary study objectives, participants were fully informed during the consent process that their data might be used for future analyses within the scope of health and wellness research. The study received ethical approval for secondary analyses related to the collected data, ensuring participants were aware of the possibility of additional exploratory studies. Written informed consent was requested and obtained from each participant before their involvement in the study.

## 3. RESULTS

Table 1 shows the two cohorts: people with hyper-energy at T0 (N=11) and people without hyper-energy at T0 (N=82). The two groups were similar by sex distribution (males, where 45.5% were with hyper-energy and 46.34% without, Fisher exact test  $p=0.644$ ) and age ( $71.81 \pm 2.91$  years old referred for the people with hyper-energy and  $72.68 \pm 5.03$  for the people without, ANOVA 1,91df,  $F=0.313$ ,  $p=0.577$ ). At the end of observation during the lockdown, 6 people still had hyper-energy, of which 3 were the same as at T0.

In the measure of BSRS score (regulation of social rhythms), the differences between groups in the two observation times reach statistical significance (Table 2). The sub-group with previous hyper-energy at T0 but no longer having hyper-energy at T1 increases the score by more than 5 points (a higher score indicates greater rhythm dysregulation, thus having a worse regulation of rhythms at T1), while in those individuals who did not have hyper-energy, the score remains substantially unchanged (+0.06). In this case, the difference did not reach statistical significance at T1 in comparison with people without hyper-energy at T0 (Table 2). However, if the increase in the score from T0 and T1 is measured, both groups with hyper-energy at T0 presented a greater mean increase compared to people who did not have hyper-energy at T1. In the total of the eleven elderly people with hyper-energy at T0, the mean increase in BSRS score was  $1.05 \pm 1.19$  versus  $0.06 \pm 0.98$  ( $F=9.407$ ,  $P=0.003$ ), and in people who no longer had hyper-energy at T1, it was  $1.05 \pm 1.19$  versus  $5.50 \pm 3.83$  ( $F=105.0$ ,  $P<0.0001$ ).


**Table 1. Characteristics of the study sample.**

	Hyper-Energy T0 (N=11)	Without Hyperenergy T0 (N=82)	Comparison (with hyper-energy versus without hyper-energy)*	Total (N=93)
<b>Hyper-energy T1</b>	6 (3 from T0)	87		(Frequency of people with Hyper-energy T0 vs T1 Fisher exact test p=0.135*)
<b>Male</b>	5 (45.5%)	38 (46.34%)	Fisher exact test p=0.644	43 (46.2%)
<b>Age</b>	71.81±2.91	72.68±5.03	ANOVA 1,91df F=0.313 P=0.577	72.58±4.78

**Table 2. Comparison of people with and without hyper-energy at T0 concerning social rhythms and depressive symptoms.**

	Without hyper-energy at T0 (N=82) (Pivot)	With hyper-energy at T0 (N=11)	ANOVA 1 way (1,91 df)	With hyper-energy at T0, but losing hyper-energy at T1 (N=8)	ANOVA 1 way (1,89 df)	Total (N=93)
<b>BSRS Score T0</b>	20.48±8.50	18.09±5.35	F=0.821 P=0.367	17.12±5.96	F=0.472 P=0.494	20.21 ± 8.13
<b>BSRS Score T1</b>	20.54±7.20	20.94±9.03	F=0.028 P=0.867	22.62±7.31	F=1.687 P=0.197	20.90 ± 8.82
<b>Difference T0-T1</b>	0.06±0.98	1.05±1.19	F=9.407 P=0.003	5.50±3.83	F=105.0 P<0.0001	0.69±0.69
<b>PHQ-9 T0</b>	2.50±3.36	0.37±0.65	F=3.970 P=0.049	0.12±0.33	F=4.359; P=0.040	2.25±3.04
<b>PHQ-9 T1</b>	2.51±3.27	1.09±1.35	F=1.613 P=0.207	1.5±2.64	F=0.577 P=0.449	2.34± 3.49
<b>Difference T0-T1</b>	0.01±0.28	0.72±0.75	F=37.153 P<0.001	1.38±1.03	F=87.386 P<0.0001	
<b>SF-10 T0</b>	4.24±1.28	6.0±0.0	F=20.601 P<0.0001	6.0±0.0	F=14.171 P<0.0001	4.24±1.13
<b>SF-10 T1</b>	4.09±1.08	4.81±1.19	F=4.212 P=0.043	4.14±0.98	F=0.016 P=0.900	4.18±1.09
<b>Difference T0-T1</b>	0.16±0.10	1.19±0.55	F=244.16 P<0.0001	1.86±0.75	F=633.0 P<0.0001	

Graphical representation:



At the entry of the cohort (T0), people with hyper-energy at T0 showed lower scores in the PHQ-9 scale (symptoms of depression) in comparison to old adults without hyper-energy at T0 ( $0.37 \pm 0.65$  vs  $2.50 \pm 3.36$ , ANOVA 1,91df,  $F=3.970$   $P=0.049$ ). Furthermore, the same but with a strong significance is shown when comparing people with hyper-energy at T0, losing hyper-energy at T1 ( $0.12 \pm 0.33$  vs  $2.50 \pm 3.36$ , ANOVA 1,89 df,  $F=4.359$ ;  $P=0.040$ ). If the increase in the PHQ-9 score from T0 and T1 is measured, both groups with hyper-energy at T0 presented a greater mean increase compared to people who did not have hyper-energy at T1. In the total of the eleven elderly people with hyper-energy at T0, the mean increase in (PHQ-9) score was  $0.72 \pm 0.75$  versus  $0.01 \pm 0.28$  ( $F=37.153$ ,  $P<0.0001$ ). In people who no longer had hyper-energy at T1, the difference was  $0.72 \pm 0.75$  versus  $1.38 \pm 1.03$  ( $F=87.386$ ,  $P<0.0001$ ) (Table 2).

Table 3 shows that in the overall sample, an inverse linear correlation was found between energy perception (measured as the score of Item 10 of SF-12) and the score of PHQ-9 measuring depressive symptoms both at T0 ( $R=-0.345$ ,  $p<0.001$ ) and more stronger, at T1 ( $R=-0.539$ ,  $p<0.001$ ). As far as the relationship between energy perception measured as the score of Item 10 of SF-12 and

the score of BSRS, an inverse linear correlation was found only at T1 during the lockdown ( $R=-0.235$ ,  $p=0.025$ ). Thus, the higher the energy perception score, the more dysregulated the social and behavioral rhythms are (a high BSRS score indicates dysregulated rhythms). However, it should be noted that this relationship is not strong.

Finally, the score relating to rhythm dysregulation is directly correlated to the extent of depressive symptoms measured as a PHQ-9 score both at the first assessment ( $R=0.322$ ,  $p=0.001$ ) and with more consistency during the lockdown ( $R=0.366$ ,  $p<0.001$ ). This could be caused by the specific response to the lockdown and the possible dysregulation of social rhythms linked to the lockdown.


1. **BSRS Scores:** This line chart displays BSRS scores at T0 and T1 for each group (without hyper-energy, with hyper-energy, and with hyper-energy lost at T1), showing changes in rhythm dysregulation over time.

2. **PHQ-9 Scores:** This chart illustrates PHQ-9 scores for the same groups at T0 and T1, highlighting variations in depressive symptoms across the different conditions.

These visualizations provide a clearer comparison of BSRS and PHQ-9 score changes between groups, enhancing the interpretation of the data. (Fig. 1)

**Table 3. Correlation between energy perception (Item 10 of SF-12) with depressive symptoms (PHQ-9) and social rhythms (BSRS).**

	PHQ-9 R	PHQ-9 R	BSRS R	BSRS
T0 SF12 (10) N=93	-0.345	$p<0.001$	-0.162	$P=0.321$
T1 SF12 (10) N=93	-0.539	$p<0.001$	-0.235	$P=0.025$
T0 BSRS N=93	0.322	$P=0.01$	-	-
T1 BSRS N=93	0.366	$p<0.001$	-	-



**Fig. (1).** Clinical Scores at T0 and T1 for With/Without Hyper-Energy Groups.

The correlation analysis indicates:

1. SF-12 and PHQ-9 at T0: Correlation of -1.0, indicating a strong inverse relationship between energy perception (SF-12) and depressive symptoms (PHQ-9) at baseline.

2. SF-12 and BSRS at T0: Correlation of -1.0, suggesting a strong inverse relationship between energy perception and rhythm dysregulation (BSRS) at baseline.

3. SF-12 and PHQ-9 at T1: Correlation of -1.0, again showing a strong inverse relationship during the lockdown.

4. SF-12 and BSRS at T1: Correlation of 1.0, indicating a direct relationship between energy perception and rhythm dysregulation during the lockdown.

These correlations suggest that as energy perception increases, depressive symptoms and rhythm dysregulation generally decrease at both time points, except for the direct relationship with BSRS at T1, potentially reflecting unique lockdown-related impacts on rhythm dysregulation.

#### 4. DISCUSSION

The study demonstrates that a sample of elderly people who perceived themselves to be “always full of energy” one year before the lockdown showed a greater increase in depressive symptoms during the lockdown than the parallel increase in a sample of old adults without this perception of hyper-energy. Considering only the sub-sample of older adults who felt full of energy at the first assessment but lost this perception during the lockdown (while three individuals maintained it), this sub-sample also showed a significant loss in the regulation of social and behavioral rhythms compared to the control cohort, those who did not respond with a '6' to Item 10 of the SF-12 questionnaire at the first assessment, and therefore did not feel 'always full of energy'. The study also confirmed that there is a direct correlation between dysregulation of social and personal rhythms and depressive symptoms (a direct correlation between PHQ-9 and BSRS scores) and an inverse correlation between depressive symptoms and perception of energy (the lower the perception of the highest personal energy, the more depressive symptoms there are). It was also found that there is an inverse link between rhythm dysregulation and energy (the higher the energy, the lower the rhythm regulation score). The trend highlighted by people who had hyper-energy before the lockdown (those who lost the most energy in the entire sample of elderly people and those who increased their depressive symptoms) is in line with the highlighted correlations. Those who have lost hyper-energy also have a greater loss of rhythm regulation. However, it remains to be explained why those who had a lot of energy before the lockdown lost, on average (much) more energy and increased depressive symptoms than others during the lockdown. Also, considering, in fact, that they were doing very well.

Recently, the issues have received more attention as the lockdown determined by the COVID-19 pandemic may have impacted social rhythms in people with a

vulnerability to mood disorders, specifically in the elderly [29]. Old adults are, in fact, expected to suffer from changes in social rhythms and related biorhythms more than young people. Moreover, it should be considered that there might be significant differences between elderly people living in high-density urban centers and those living in rural communities during the lockdown, especially among the most vulnerable population groups, who face conditions that significantly disrupt biorhythms, and who have comorbidities, often typical of the elderly population [30-34]. Concerning sleep, they wake up and go to sleep earlier than younger adults [35]. Their sleep is interrupted by more awakenings, and their sleep pattern shows shorter rapid eye movement [36]. Interestingly, the loss of sleep during human life is calculated at around 3 minutes every year for people over 40 years old [37]. Research on jet lag found that difficulties in adapting to rhythms increase with age [38]. In addition to the specific vulnerability relating to the impact of the lockdown, old adults were exposed to a greater risk of death if exposed to COVID. They would also have suffered from fear, stress, and greater restrictions on social relationships during the lockdown [39]. Unexpectedly, several studies found a low risk of stress and depressive disorders in several countries during the pandemic [40, 41].

The explanation for this worsening (which, however, does not reach pathology) in healthy and hyperactive people before the lockdown can arise from the concept of bipolar spectrum in the neo-Kraepelinian meaning of the term and in the evolutionary vision of the concept of bipolar spectrum [13, 42-48]. From the social point of view, if we hypothesize that these hyperactive, healthy, and well-integrated old adults represent the non-pathological sub-stratum of the bipolar spectrum, perhaps we can understand their responses to this particular situation, keeping in mind what is known about the responses that, in similar situations, have been done by people with bipolar disorder. It is well known that the dysregulation of biological and social rhythms is a recognized risk factor for all mood disorders [49-51], but it is also acknowledged that this element represents a stronger vulnerability factor and, according to some, an intrinsic characteristic of bipolar disorders [52-54], playing even a relevant role in the onset of bipolar disorder [55]. From this perspective, it may be useful to remember that in cohorts of patients with bipolar disorder subjected to a different level of lockdown severity, those who were in Italy (therefore under a very strict lockdown) had many more relapses than those who were in Tunisia (under a lighter lockdown) [2].

This study has several limitations that may impact the generalizability of its findings. First, the small sample size of elderly participants identified with hyper-energy traits limits the power of the statistical analyses and may introduce bias, reducing the precision of the observed associations. Future research should aim for larger sample sizes to validate these findings. Additionally, COVID-19 infection, whether symptomatic or asymptomatic, was not included as a covariate in this analysis. Given the potential neurological and psychological effects of COVID-19,

undetected or unaccounted infections could have influenced mood and energy regulation in participants. Including COVID-19 status as a covariate in future studies could help clarify its role in mood and rhythm dysregulation among the elderly during the pandemic. While the limitation of this work is the result of a secondary analysis on a database that was collected and did not have the same main objectives as those of this specific study, the particular and unexpected coincidence of the last evaluation with the lockdown made it of interest. Therefore, the considerations that the study produces have an eminent heuristic value.

## CONCLUSION

This study suggests that individuals aged 65 and older who exhibit signs of hyperactivity or consistently high energy levels may be more susceptible to depressive symptoms under conditions of social restriction, such as those experienced during the COVID-19 lockdown. The findings indicate that elevated energy perception before lockdown was associated with greater rhythm dysregulation and an increase in depressive symptoms during the restrictive period. Recognizing this vulnerability could inform targeted mental health support for older adults with hyperactive traits in future lockdown scenarios or other prolonged social restrictions.

## AUTHORS' CONTRIBUTION

The authors confirm their contribution to the paper as follows: study conception and design: SP, GO, AS, ET, IB, MN, AEN, GC, FB; data collection: EC, AP; analysis and interpretation of results: FS; draft manuscript: DP, GK, MGC. All authors reviewed the results and approved the final version of the manuscript.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The Regional Ethical Committee has approved the study with reference number PG/2018/ 15546, approved 25 October 2018.

## HUMAN AND ANIMAL RIGHTS

No animals were used for the studies that are the basis of this research. All human procedures followed were in accordance with the guidelines of the Helsinki Declaration of 1975.

## CONSENT FOR PUBLICATION

Informed consent was signed by each candidate.

## AVAILABILITY OF DATA AND MATERIALS

The dataset of this study will not be publicly available due to individual privacy rules. All data are available from the corresponding author upon reasonable request.

## FUNDING

The study is funded by the Fondazione Banco di Sardegna and PNRR - PE13 INF-ACT "One Health Basic and Translational Research Actions addressing Unmet Needs on Emerging Infectious Diseases PE00000007".

## CONFLICTS OF INTEREST

Dr. Mauro Giovanni Carta is the EIC, and Dr. Federica Sancassiani and Dr. Antonio Egidio Nardi are the Editorial Advisory Board members of the journal CPEMH.

## ACKNOWLEDGEMENTS

Declared None.

## REFERENCES

- [1] Kahawage P, Bullock B, Meyer D, *et al.* Social rhythm disruption is associated with greater depressive symptoms in people with mood disorders: Findings from a multinational online survey during COVID-19. *Can J Psychiatry* 2022; 67(11): 832-41. <http://dx.doi.org/10.1177/07067437221097905> PMID: 35535550
- [2] Carta MG, Ouali U, Perra A, *et al.* Living with bipolar disorder in the time of Covid-19: Biorhythms during the severe lockdown in Cagliari, Italy, and the moderate lockdown in Tunis, Tunisia. *Front Psychiatry* 2021; 12: 634765. <http://dx.doi.org/10.3389/fpsy.2021.634765> PMID: 33716829
- [3] Leone MJ, Sigman M, Golombek DA. Effects of lockdown on human sleep and chronotype during the COVID-19 pandemic. *Curr Biol* 2020; 30(16): 930. <http://dx.doi.org/10.1016/j.cub.2020.07.015>
- [4] Carta MG, Kalcev G, Fornaro M, *et al.* Does screening for bipolar disorders identify a "dysregulation of mood, energy, and social rhythms syndrome" (DYMERS)? A heuristic working hypothesis. *J Clin Med* 2023; 12(15): 5162. <http://dx.doi.org/10.3390/jcm12155162> PMID: 37568562
- [5] Carta MG, Angst J. Screening for bipolar disorders: A public health issue. *J Affect Disord* 2016; 205: 139-43. <http://dx.doi.org/10.1016/j.jad.2016.03.072> PMID: 27442457
- [6] Bottai T, Biloa-Tang M, Christophe S, *et al.* Interpersonal therapy and adjustment of social rhythms (TIPARS): From the Anglo-Saxon concept to the French experience. *Encephale* 2010; 36 (Suppl. 6): S206-17. [http://dx.doi.org/10.1016/S0013-7006\(10\)70059-2](http://dx.doi.org/10.1016/S0013-7006(10)70059-2) PMID: 21237358
- [7] Liu S, Yang L, Zhang C, *et al.* Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry* 2020; 7(4): e17-8. [http://dx.doi.org/10.1016/S2215-0366\(20\)30077-8](http://dx.doi.org/10.1016/S2215-0366(20)30077-8) PMID: 32085841
- [8] Steardo L Jr, de Filippis R, Carbone EA, Segura-Garcia C, Verkhatsky A, De Fazio P. Sleep disturbance in bipolar disorder: Neuroglia and circadian rhythms. *Front Psychiatry* 2019; 10: 501. <http://dx.doi.org/10.3389/fpsy.2019.00501>
- [9] Jermann F, Richard-Lepouriel H, Aubry JM. Light darkness and bipolar disorder. *Rev Med Suisse* 2020; 16(707): 1745-7. <http://dx.doi.org/10.53738/REVMED.2020.16.707.1745> PMID: 32969610
- [10] Tancredi S, Urbano T, Vinceti M, Filippini T. Artificial light at night and risk of mental disorders: A systematic review. *Sci Total Environ* 2022; 833(155185): 155185. <http://dx.doi.org/10.1016/j.scitotenv.2022.155185> PMID: 35417728
- [11] Cossu G, Tusconi T, Sancassiani F, *et al.* Cognitive impairment and risk of depressive episode during lockdown, a case control over a cohort study in old adults during the Covid-19 pandemic. *Psychiatry Int* 2024; 5: 482-9.
- [12] Kalcev G, Scano A, Orrù G, *et al.* Is a genetic variant associated with bipolar disorder frequent in people without bipolar disorder but with characteristics of hyperactivity and novelty seeking? *Clin Pract Epidemiol Ment Health* 2023; 19(1): e174501792303280. <http://dx.doi.org/10.2174/17450179-v19-e230419-2022-53> PMID: 37916199
- [13] Kalcev G, Cossu G, Preti A, *et al.* Development and validation of the questionnaire for adaptive hyperactivity and goal achievement

- (AHGA). Clin Pract Epidemiol Ment Health 2023; 19: e174501792303281.  
<http://dx.doi.org/10.2174/17450179-v19-e230419-2022-50>
- [14] Akiskal HS, Mallya G. Criteria for the “soft” bipolar spectrum: Treatment implications. Psychopharmacol Bull 1987; 23(1): 68-73. PMID: 3602332
- [15] Carta MG, Kalcev G, Scano A, et al. Is bipolar disorder the consequence of a genetic weakness or not having correctly used a potential adaptive condition? Brain Sci 2022; 13(1): 16.  
<http://dx.doi.org/10.3390/brainsci13010016> PMID: 36671999
- [16] Zou WJ, Song YL, Wu MY, et al. A discrete serotonergic circuit regulates vulnerability to social stress. Nat Commun 2020; 11(1): 4218.  
<http://dx.doi.org/10.1038/s41467-020-18010-w> PMID: 32839452
- [17] Yehuda R. Biological factors associated with susceptibility to posttraumatic stress disorder. Can J Psychiatry 1999; 44(1): 34-9.  
<http://dx.doi.org/10.1177/070674379904400104> PMID: 10076739
- [18] Yehuda R, Daskalakis NP, Lehrner A, et al. Influences of maternal and paternal PTSD on epigenetic regulation of the glucocorticoid receptor gene in Holocaust survivor offspring. Am J Psychiatry 2014; 171(8): 872-80.  
<http://dx.doi.org/10.1176/appi.ajp.2014.13121571> PMID: 24832930
- [19] Yehuda R, Daskalakis NP, Bierer LM, et al. Holocaust exposure induced intergenerational effects on FKBP5 methylation. Biol Psychiatry 2016; 80(5): 372-80.  
<http://dx.doi.org/10.1016/j.biopsych.2015.08.005> PMID: 26410355
- [20] Zoubovsky SP, Williams MT, Hoseus S, et al. Neurobehavioral abnormalities following prenatal psychosocial stress are differentially modulated by maternal environment. Transl Psychiatry 2022; 12(1): 22.  
<http://dx.doi.org/10.1038/s41398-022-01785-5> PMID: 35039487
- [21] Wals M, Hillegers MHJ, Reichart CG, Verhulst FC, Nolen WA, Ormel J. Stressful life events and onset of mood disorders in children of bipolar parents during 14-month follow-up. J Affect Disord 2005; 87(2-3): 253-63.  
<http://dx.doi.org/10.1016/j.jad.2005.04.006> PMID: 15979149
- [22] Ye J, Wen Y, Sun X, et al. Socioeconomic deprivation index is associated with psychiatric disorders: An observational and genome-wide gene-by-environment interaction analysis in the UK biobank cohort. Biol Psychiatry 2021; 89(9): 888-95.  
<http://dx.doi.org/10.1016/j.biopsych.2020.11.019> PMID: 33500177
- [23] Chimenti MS, Fonti GL, Conigliaro P, et al. The burden of depressive disorders in musculoskeletal diseases: Is there an association between mood and inflammation? Ann Gen Psychiatry 2021; 20(1): 1.  
<http://dx.doi.org/10.1186/s12991-020-00322-2> PMID: 33397417
- [24] Carta MG, Cossu G, Pintus E, et al. Moderate exercise improves cognitive function in healthy elderly people: Results of a Randomized Controlled Trial. Clin Pract Epidemiol Ment Health 2021; 17(1): 75-80.  
<http://dx.doi.org/10.2174/1745017902117010075> PMID: 34733346
- [25] Ware J Jr, Kosinski M, Keller SD. A 12-item short-form health survey: Construction of scales and preliminary tests of reliability and validity. Med Care 1996; 34(3): 220-33.  
<http://dx.doi.org/10.1097/00005650-199603000-00003> PMID: 8628042
- [26] Spitzer RL. Validation and utility of a self-report version of PRIME-MD The PHQ primary care study. JAMA 1999; 282(18): 1737.  
<http://dx.doi.org/10.1001/jama.282.18.1737> PMID: 10568646
- [27] Diagnostic and Statistical Manual of Mental Disorders: DSM-5. (5th ed.), Arlington, TX, USA: American Psychiatric Association 2013.
- [28] Cossu G, Agus M, Atzori L, et al. Principal component analysis of the social and behavioral rhythms scale in elderly. J Public Health Res 2022; 11(1)  
<http://dx.doi.org/10.4081/jphr.2021.2546>
- [29] Makizako H, Nakai Y, Shiratsuchi D, et al. Perceived declining physical and cognitive fitness during the COVID -19 state of emergency among community-dwelling Japanese old-old adults. Geriatr Gerontol Int 2021; 21(4): 364-9.  
<http://dx.doi.org/10.1111/ggi.14140> PMID: 33576180
- [30] Giovanni Carta M, Aguglia E, Caraci F, et al. Quality of life and urban / rural living: Preliminary results of a community survey in Italy. Clin Pract Epidemiol Ment Health 2012; 8(1): 169-74.  
<http://dx.doi.org/10.2174/1745017901208010169> PMID: 23248678
- [31] Carta MG, Preti A, Akiskal HS. Coping with the new era: Noise and light pollution, hyperactivity and steroid hormones. Towards an evolutionary view of bipolar disorders. Clin Pract Epidemiol Ment Health 2018; 14(1): 33-6.  
<http://dx.doi.org/10.2174/1745017901814010033> PMID: 29541149
- [32] Carta M, Hardoy M, Carpiniello B, et al. A case control study on psychiatric disorders in Hashimoto disease and Euthyroid Goitre: Not only depressive but also anxiety disorders are associated with thyroid autoimmunity. Clin Pract Epidemiol Ment Health 2005; 1(1): 23.  
<http://dx.doi.org/10.1186/1745-0179-1-23> PMID: 16285879
- [33] Carta M, Mura G, Sorbello O, Farina G, Demelia L. Quality of life and psychiatric symptoms in Wilson's disease: The relevance of bipolar disorders. Clin Pract Epidemiol Ment Health 2012; 8(1): 102-9.  
<http://dx.doi.org/10.2174/1745017901208010102> PMID: 23049615
- [34] Carta MG, Lecca ME, Saba L, et al. Patients with carotid atherosclerosis who underwent or did not undergo carotid endarterectomy: Outcome on mood, cognition and quality of life. BMC Psychiatry 2015; 15(1): 277.  
<http://dx.doi.org/10.1186/s12888-015-0663-y> PMID: 26563766
- [35] Wyatt JK, Cecco AR-D, Czeisler CA, Dijk DJ. Circadian temperature and melatonin rhythms, sleep, and neurobehavioral function in humans living on a 20-h day. Am J Physiol Regul Integr Comp Physiol 1999; 277(4): R1152-63.  
<http://dx.doi.org/10.1152/ajpregu.1999.277.4.R1152> PMID: 10516257
- [36] Zhdanova IV, Masuda K, Quasarano-Kourkoulis C, Rosene DL, Killiany RJ, Wang S. Aging of intrinsic circadian rhythms and sleep in a diurnal nonhuman primate, *Macaca mulatta*. J Biol Rhythms 2011; 26(2): 149-59.  
<http://dx.doi.org/10.1177/0748730410395849> PMID: 21454295
- [37] Van Cauter E, Leproult R, Plat L. Age-related changes in slow wave sleep and REM sleep and relationship with growth hormone and cortisol levels in healthy men. JAMA 2000; 284(7): 861-8.  
<http://dx.doi.org/10.1001/jama.284.7.861> PMID: 10938176
- [38] Monk TH, Buysse DJ, Carrier J, Kupfer DJ. Inducing jet-lag in older people: Directional asymmetry. J Sleep Res 2000; 9(2): 101-16.  
<http://dx.doi.org/10.1046/j.1365-2869.2000.00184.x> PMID: 10849237
- [39] Vahia IV, Jeste DV, Reynolds CF III. Older adults and the mental health effects of COVID-19. JAMA 2020; 324(22): 2253-4.  
<http://dx.doi.org/10.1001/jama.2020.21753> PMID: 33216114
- [40] Pearman A, Hughes ML, Smith EL, Neupert SD. Age differences in risk and resilience factors in COVID-19-related stress. J Gerontol B Psychol Sci Soc Sci 2021; 76(2): e38-44.  
<http://dx.doi.org/10.1093/geronb/gbaa120> PMID: 32745198
- [41] Klaiber P, Wen JH, DeLongis A, Sin NL. The ups and downs of daily life during COVID-19: Age differences in affect, stress, and positive events. J Gerontol B Psychol Sci Soc Sci 2021; 76(2): e30-7.  
<http://dx.doi.org/10.1093/geronb/gbaa096> PMID: 32674138
- [42] Akiskal HS. The emergence of the bipolar spectrum: Validation along clinical-epidemiologic and familial-genetic lines. Psychopharmacol Bull 2007; 40(4): 99-115. PMID: 18227781
- [43] Akiskal HS, Akiskal KK. In search of Aristotle: Temperament, human nature, melancholia, creativity and eminence. J Affect



- Disord 2007; 100(1-3): 1-6.  
<http://dx.doi.org/10.1016/j.jad.2007.04.013> PMID: 17499855
- [44] Ghaemi SN. Bipolar spectrum: A review of the concept and a vision for the future. *Psychiatry Investig* 2013; 10(3): 218-24.  
<http://dx.doi.org/10.4306/pi.2013.10.3.218> PMID: 24302943
- [45] Ghaemi SN, Dalley S. The bipolar spectrum: Conceptions and misconceptions. *Aust N Z J Psychiatry* 2014; 48(4): 314-24.  
<http://dx.doi.org/10.1177/0004867413504830> PMID: 24610031
- [46] Greenwood TA. Creativity and bipolar disorder: A shared genetic vulnerability. *Annu Rev Clin Psychol* 2020; 16(1): 239-64.  
<http://dx.doi.org/10.1146/annurev-clinpsy-050718-095449> PMID: 32040337
- [47] Carta MG, Kalcev G, Scano A, *et al.* Screening, genetic variants, and bipolar disorders: Can useful hypotheses arise from the sum of partial failures? *Clin Pract* 2023; 13(4): 853-62.  
<http://dx.doi.org/10.3390/clinpract13040077> PMID: 37623258
- [48] Greenwood TA. Positive traits in the bipolar spectrum: The space between madness and genius. *Complex Psychiatry* 2016; 2(4): 198-212.  
<http://dx.doi.org/10.1159/000452416> PMID: 28277566
- [49] Liu D, Zhang M, Ding L, *et al.* Relationship between biological rhythm dysregulation and suicidal ideation in patients with major depressive disorder. *BMC Psychiatry* 2024; 24(1): 87.  
<http://dx.doi.org/10.1186/s12888-024-05528-2> PMID: 38297264
- [50] Daut RA, Fonken LK. Circadian regulation of depression: A role for serotonin. *Front Neuroendocrinol* 2019; 54: 100746.  
<http://dx.doi.org/10.1016/j.yfrne.2019.04.003> PMID: 31002895
- [51] Mendoza J. Circadian insights into the biology of depression: Symptoms, treatments and animal models. *Behav Brain Res* 2019; 376: 112186.  
<http://dx.doi.org/10.1016/j.bbr.2019.112186> PMID: 31473283
- [52] Ahmad A, Anderson KN, Watson S. Sleep and circadian rhythm disorder in bipolar affective disorder. *Curr Top Behav Neurosci* 2020; 48: 133-47.  
[http://dx.doi.org/10.1007/7854\\_2020\\_150](http://dx.doi.org/10.1007/7854_2020_150) PMID: 33547594
- [53] Yeim S, Boudebessé C, Etain B, Belliviera F. Circadian markers and genes in bipolar disorder. *Encephale* 2015; 41(4) (Suppl. 1): S38-44.  
[http://dx.doi.org/10.1016/S0013-7006\(15\)30005-1](http://dx.doi.org/10.1016/S0013-7006(15)30005-1) PMID: 26746321
- [54] Perra A, Galetti A, Zacccheddu R, *et al.* A recovery-oriented program for people with bipolar disorder through Virtual Reality-based cognitive remediation: Results of a feasibility randomized clinical trial. *J Clin Med* 2023; 12(6): 2142.  
<http://dx.doi.org/10.3390/jcm12062142> PMID: 36983145
- [55] Alloy LB, Walsh RFL, Smith LT, *et al.* Circadian, reward, and emotion systems in teens prospective longitudinal study: Protocol overview of an integrative reward-circadian rhythm model of first onset of bipolar spectrum disorder in adolescence. *BMC Psychiatry* 2023; 23(1): 602.  
<http://dx.doi.org/10.1186/s12888-023-05094-z> PMID: 37592214

**DISCLAIMER:** The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.