


Clinical Practice & Epidemiology in Mental Health

Content list available at: <https://clinical-practice-and-epidemiology-in-mental-health.com>



RESEARCH ARTICLE

Validity and Reliability of the Digital Functioning Assessment Short Test (D-FAST) in the Brazilian Sample

Silvia Dubou Serafim^{1,2}, Jeferson Ferraz Goularte^{1,2}, Marco Antonio Caldieraro^{1,2}, Flavia Moreira Lima³, Giovana Dalpiaz^{1,2}, Francisco Diego Rabelo-da-Ponte^{1,2}, Carla Torrent⁴, Brisa Solé⁴, Eduard Vieta⁴ and Adriane Ribeiro Rosa^{1,2,5,*}

¹Laboratory of Molecular Psychiatry, Hospital de Clínicas de Porto Alegre, University Federal of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, RS, Brazil

²Postgraduate Program in Psychiatry and Behavioral Sciences, University Federal of Rio Grande do Sul, Porto Alegre, RS, Brazil

³Hospital de Clínicas de Porto Alegre, University Federal of Rio Grande do Sul, Porto Alegre, Rio Grande do Sul, RS, Brazil

⁴Bipolar and Depressive Disorders Unit, Hospital Clinic, University of Barcelona, Institute of Neuroscience, IDIBAPS, CIBERSAM, c/Villarreal, 170, 12-0, 08036 Barcelona, Catalonia, Spain

⁵Department of Pharmacology, Institute of Basic Health Sciences, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil

Abstract:

Background:

The COVID-19 pandemic has caused major disruptions in all aspects of daily functioning, from school and work to interactions with friends and family. The Functioning Assessment Short Test (FAST) is an interviewer-administered scale validated in the psychiatric sample with no previous study assessing its validity and reliability in a digital format. Thus, we aimed to analyse the psychometric properties of the digital version of the FAST and understand the implications of COVID-19 and restrictive measures on functioning.

Methods:

Data were collected using an online survey. The psychometric properties of the digital FAST were assessed by confirmatory factor analysis, Cronbach's alpha, and discriminant functional by cluster analysis in a community sample.

Results:

Out of the total sample, 2,543 (84.1%) were female, and the mean (SD) age was 34.28 (12.46) years. The digital FAST retained the six factors structure observed in the original version, with Cronbach's alpha above 0.9. In addition, we showed evidence of discriminant validity by differentiating three clusters of psychosocial functioning. Clinical and demographic differences between groups explained, in part, the heterogeneity of functioning, thus providing support for the construct validity of the instrument.

Conclusion:

The digital FAST is a simple and easy-to-understand instrument that provides a multidimensional assessment of functioning without the need for an interviewer. Furthermore, our findings may help to better understand the psychosocial implications of the pandemic and the importance of planning specific interventions to rehabilitate the affected group.

Keywords: COVID-19, Pandemic, Public health, Deaths, Virus transmission, Psychosocial implications.

Article History

Received: April 11, 2022

Revised: September 2, 2022

Accepted: September 7, 2022

1. INTRODUCTION

On January 30th, 2020, the World Health Organization (WHO) announced that severe acute respiratory syndrome coronavirus (COVID-19) was a public health emergency of international concern. Currently, COVID-19 has infected over 125 million people and resulted in over two million deaths

worldwide, while in Brazil, it has infected over 21 million people and resulted in over 610 thousand deaths (as of November 13th, 2021) [1]. According to the WHO, respiratory etiquette, hand washing, and physical distancing must be maintained to control virus transmission, while most people have not been vaccinated. However, the pandemic and restrictive measures may be particularly harmful to low-income and middle-income countries in which weak social safety nets and economic resources are not sufficient to cover daily needs

* Address correspondence to this author at the Postgraduate Program in Psychiatry and Behavioral Sciences, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil; E-mail: adriannerosa@gmail.com

[2], such as Brazil, where the greatest risk of disease transmission is among the poorest communities [3].

The pandemic and containment measures [4] have changed daily routines, bringing several physical and mental consequences, such as emotional distress and increased risk of psychiatric illnesses, especially among the most vulnerable groups [5]. Indeed, mental health burdens and increased use of mental health services are expected as a consequence of the pandemic. The high prevalence of anxiety, depression, and other stress-related disorders observed during the acute phase of COVID-19 has been consistently reported in multicultural studies [6, 5, 7 - 9]. Notably, these psychiatric conditions are among the leading contributors to disability worldwide [10]. Thus, the relationship between mental health and functional impairment is not new in psychiatry.

Psychosocial functioning describes a person's ability to perform the tasks of everyday life, engage in relationships with others in ways that are gratifying to the individual and others, and meet the needs of the community in which the person lives [11]. The COVID-19 pandemic caused major disruptions in all aspects of functioning, from school and work to interactions with friends or family and recreational time. Furthermore, the measures to mitigate the disease have substantially altered the economic scenario with an increase in unemployment and uncertainty about the future [12], which contribute to the worsening of mental health. Thus, a better understanding of the pattern of psychosocial functioning among general populations during the COVID-19 pandemic would be of clinical utility as such information could contribute to the development of interventions focused on functional restoration.

The Functioning Assessment Short Test (FAST) [13] is an interviewer-administered scale involving the use of pen and paper that allows the multidimensional evaluation of functioning, including aspects, such as the individual's ability to function socially or occupationally or to live independently as well as financial issues and cognition. The FAST scale was originally validated in several languages in distinct clinical samples [13 - 21] as well as in healthy individuals [22, 23], with no previous study assessing its validity in self-reported digital format. The issue with transferring such validated instruments to the digital format has been raised by some authors [24], with others suggesting that a validated pen and paper scale needs further validation when used in other formats, including online assessments [25]. In addition, although many researchers have assessed the impacts of COVID-19 on mental health [6, 5, 7 - 9], no data are yet available on the effects of the current pandemic on psychosocial functioning.

Hence, the purpose of the present study was to analyze the psychometric properties of the online self-reported version of the FAST and also to understand the implications of COVID-19 and restrictive measures on psychosocial functioning by cluster analyses in a subsample of the Brazilian population during the first wave of coronavirus transmission.

2. MATERIALS AND METHODS

2.1. Participants

A cross-sectional web-based survey was carried out using an anonymous online questionnaire distributed via social networks. The data were collected from May 20th to September 13th, 2020, during the first peak of COVID-19 in Brazil. The online questionnaire consisted of sociodemographic variables and the assessment of psychosocial functioning, physical, and mental health status, including the history of previous psychiatric disorders and the severity of stress, anxiety, and depression as described below. The inclusion criteria included being older than 18 and living in Brazil at the time of the survey. All participants provided online informed consent. The local ethical committees approved all procedures.

2.2. Assessments

a) Psychosocial functioning was assessed by the Functioning Assessment Short Test (FAST) scale [13], which allows for the evaluation of the main aspects of functioning: autonomy, occupational functioning, cognitive functioning, interpersonal relationships, financial issues, and leisure time. All items of scale are rated using a four-point Likert scale: 0 = no difficulty, 1 = mild difficulty, 2 = moderate difficulty, and 3 = severe difficulty. The global score is the sum of all items. The higher the score, the more severe the difficulties. The FAST is an interviewer-administered, transdiagnostic scale validated in distinct clinical samples and available in several languages [13 - 21]. It was also validated in healthy individuals [22, 23]. In this web-based survey, the online self-reported version of the FAST was used for the first time; a brief description of each item was included in this version in order to guarantee the best understanding by the responders (supplementary material).

b) The Impact of Event Scale-Revised (IES-R) is a self-rated 22-item questionnaire divided into three domains (avoidance, intrusion, and hyperarousal), which evaluates the degree of distress caused by a traumatic event [26]. Each item is rated on a five-point Likert scale (0 = not at all; 1 = a little bit; 2 = moderately; 3 = quite a bit; 4 = extremely). The IES-R total score is the sum of the average of each domain. A total score higher than 5.6 indicates psychological stress.

c) The Patient-Reported Outcomes Measurement Information System (PROMIS) for depression (PROMIS Short Form v1.0 - Depression 8a) assesses negative mood (sadness, guilt), views of self (self-criticism, worthlessness), social cognition (loneliness, interpersonal alienation), and decreased positive affect and engagement (loss of interest, meaning, and purpose).

d) The PROMIS anxiety assesses self-reported fear (fearfulness, panic), anxious misery (worry, dread), hyperarousal (tension, nervousness, restlessness), and somatic symptoms related to arousal (racing heart, dizziness).

Both PROMIS instruments consist of an eight-item questionnaire that assesses symptoms over the period of seven days, with items rated on a five-point Likert scale (1 = never; 2 = rarely; 3 = sometimes; 4 = often; 5 = always). All PROMIS

scores were presented as T-scores calculated by the Health Measures Scoring Service (https://www.assessmentcenter.net/ac_scoringservice) from the raw sum score, using T-scores from the United States general population. The T-score is the standardized score with a mean of 50 and a standard deviation of 10. For depression and anxiety, T-scores lower or equal to 55 indicate no significant symptoms, higher than 55 to 60 indicate mild symptoms, higher than 60 to 70 indicate moderate symptoms, and higher than 70 to 83.1 and 81.1, respectively, indicate severe symptoms.

3. STATISTICAL ANALYSIS

3.1. Confirmatory Factor Analysis

R (version 4.0.2) and RStudio (version 3.5.3) software were used for all analyses. We applied a confirmatory factor analysis (CFA) to identify the factorial structures of a set of items. Furthermore, CFA is highly useful in verifying the relationship between observed variables and latent traits. We performed a CFA through the principal axis factoring method in order to describe the internal structure of the online self-reported FAST scale and to confirm the same factors found in the pen and paper version of FAST using the package “lavaan” (version 0.6-12). Then, we used the oblimin rotation with the Kaiser-Meyer-Olkin (KMO) measure of sampling adequacy (>0.5) and Bartlett's Test of Sphericity ($p < 0.05$) to confirm whether those metrics met all assumptions for CFA [27]. Afterward, we investigated all eigenvalues over Kaiser's criterion of 1 to confirm the number of factors and the number of items for each factor.

3.2. Internal Reliability

The Cronbach's alpha was used to analyze the internal reliability of the online FAST global factor (*i.e.*, FAST total score) and was assessed using the following criteria: $\alpha \leq 0.9$, excellent; $\alpha \leq 0.8$, good; $\alpha \leq 0.7$, adequate (package “ltm”, version 1.2-0) [28]. Cronbach's alpha is a coefficient of reliability among raters; in other words, a high value for Chronbach's alpha means all the psychometric items measure the same construct.

3.3. Discriminant Validity

All functioning domains were converted into Z-score using all samples. Afterward, we performed the Partition Around Medoids (PAM) algorithm [29] to identify functioning clusters of subjects during the COVID-19 outbreak (package *fpc*, version 2.2-9) as a proxy of discriminant validity. PAM algorithm was used rather than k-means, a classical clustering algorithm, because it is more robust to noise and outliers, minimizing the sum of dissimilarities between data points. To calculate the dissimilarities between pairs of subjects, Gower's distance was applied. The optimal number of clusters was determined by the Gap statistic method using the package “factoextra” (version 1.0.7). After the clustering, a discriminant function analysis (DFA) was performed using the package “MASS” (version 7.3-51.6) to confirm the clusters retained and investigate the predictive power of the clustering of each individual's functioning domain to the functioning subgroup.

3.4. Univariate and Multivariate Analysis

Furthermore, we conducted comparisons (one-way ANOVA with Tukey HSD post-hoc test and χ^2 applied as appropriate) between the different clusters to examine possible differences in functional status, demographic and clinical variables. Moreover, multinomial logistic regression (package “nnet,” version 7.3-14) was carried out using the good functioning cluster as the reference category outcome in order to identify predictors of functioning. The model included the following variables: age, T-score of depression, T-score of anxiety, gender, work status, household income, work status in the ongoing pandemic, previous history of psychiatric illness, marital status, education, and post-traumatic stress. We also calculated the effect size using Hedges' *g* from the mean and standard deviation between good functioning vs. intermediate functioning and good functioning vs. low functioning (package “esc,” version 0.5.1). Positive values for Hedges' *g* mean a good functioning cluster that shows better performance than intermediate or poor functioning. Statistical significance was set at $p < .05$.

4. RESULTS

4.1. Demographic Characteristics

A total of 3,023 individuals completed the survey. Out of the total sample, 2,543 (84.1%) were female, and the mean (SD) age was 34.28 (12.46) years.

4.2. Confirmatory Factor Analysis

The original version of FAST is based on a six-factor structure [13]. In this study, the online self-reported FAST was assessed for fitting the same structure. Initially, the Kaiser-Meyer-Olkin test verified the sampling adequacy for the analysis (KMO = 0.91) with good values for Bartlett's test for sphericity ($\chi^2(276) = 32,288$, $p < 0.001$). In addition, an initial analysis was performed to obtain eigenvalues for each factor in the data. In agreement with the original FAST, the online FAST showed that six factors had eigenvalues over Kaiser's criterion of 1 and explained 63.08% of the variance in combination. We retained six factors due to the large sample size and the convergence of the scree plot and Kaiser's criterion. The same internal structure (*i.e.*, six factors) of the original version was also reported in the online self-reported FAST except for two items, *i.e.*, “participating in social activities” and “having satisfactory sexual relationships,” shifting from the interpersonal relationships factor to the leisure time factor. However, as leisure time and interpersonal relationships were strongly related, this change did not compromise the internal structure of the online FAST. Hence, the maximum score of the interpersonal relationships and leisure time domains was 12 in the online version compared to scores of 18 and 6, respectively, as described in the original scale. The internal consistency coefficient presented an overall Cronbach = 0.95, indicating excellent internal reliability of the online FAST.

4.3. Discriminant Validity

The PAM algorithm, through the Gap statistic method,

provided evidence for three functional clusters among the 3,023 volunteers, as demonstrated in Fig. (1). The first cluster included 661 subjects (21.86%) who presented low functioning (LF). The second one contained 1,436 subjects (47.50%) with intermediate functioning results (IF). The last cluster had 926 patients (30.36%) with good functioning (GF). The discriminant function analysis (DFA) exhibited two discriminant functions, which explained 99.4% and 0.6% of the variance, respectively (Wilks' $\lambda = 0.19$, $\chi^2(12) = 4884.41$, $p < 0.001$; Wilks' $\lambda = 0.97$, $\chi^2(5) = 65.1$, $p < 0.001$). The subjects were classified by DFA into 89.2% of the cases, demonstrating the validity of the three functioning clusters. The cognitive domain and leisure time domain were among the domains that better classified participants into functioning clusters (Function

1: $r = 0.46$ and $r = -0.37$; function 2: $r = 0.4$ and $r = 0.67$, respectively) Fig. (2) for graphical agglomeration of the functional subgroups).

The LF group showed marked impairment in all FAST subdomains, with a huge difference in mean scores between LF and GF. The IF cluster exhibited an intermediate level of functioning in all subdomains, with a great difference in mean scores between IF and GF. Finally, the GF cluster presented a high-functioning performance in all subdomains (Table 1). Taken together, these data suggested that the online self-reported FAST scale could discriminate subjects with different levels of psychosocial functioning, thus supporting the discriminant validity.




Fig. (1). Mean of functioning performance between three clusters of individuals during COVID-19 pandemic. The X-axis is the functioning domains and Y-axis is the value of z-score based on mean and standard deviation of all sample.




Fig. (2). Discriminant validity: graphical agglomeration of the functional subgroups.

Table 1. Comparisons between the three functional clusters across overall and specific functioning domains of the FAST using one-way ANOVA with Tukey HSD post hoc test. Hedge’s g for a measure of effect size.

	Low Functioning n= 661 Mean(SD)	Intermediate Functioning n=1436 Mean (SD)	Good Functioning n=926 Mean (SD)	F-statistics	p-value	Hedge’s g (95%CI) Good vs. Low	Hedge’s g Good vs. intermediate
Autonomy	6.01(2.60) ^a	3.14 (2.03) ^b	0.92 (1.41) ^c	1233.29	<.001	-2.5517 (-2.6853; -2.4181)	-1.2245 (-1.3142; -1.1348)
Occupational	9.70 (3.60) ^a	6.05 (3.33) ^b	2.19 (2.46) ^c	1106.60	<.001	-2.5122 (-2.6449;-2.3795)	-1.2782 (-1.3684;-1.1879)
Financial	3.94 (1.68) ^a	2.11 (1.46) ^b	0.67 (1.11) ^c	1034.82	<.001	-2.3747 (-2.5043;-2.2452)	-1.0793 (-1.1674; -0.9911)
Cognitive	8.98 (3.11) ^a	5.15 (2.58) ^b	2.29 (2.08) ^c	1305.18	<.001	-2.6123 (-2.7473;-2.4773)	-1.1930 (-1.2824; -1.1037)
Interpersonal	6.76 (2.68) ^a	3.67 (2.22) ^b	1.58 (1.82) ^c	1217.09	<.001	-2.3333 (-2.4620;-2.2046)	-1.0082 (-1.0956;-0.9207)
Leisure time	9.66 (1.93) ^a	6.89 (2.29) ^b	3.98 (2.44) ^c	981.95	<.001	-2.5325 (-2.6657; -2.3994)	-1.2379 (-1.3278; -1.1481)
FAST total	45.07 (8.05) ^a	27.00 (6.31) ^b	11.64 (5.61) ^c	5064.70	<.001	-4.9618 (-5.1612;-4.7623)	-2.5400 (-2.6499; -2.4301)

Note: Different letters mean difference between clusters: aLow Functioning vs. Intermediate Functioning; bIntermediate Functioning vs. Good Functioning; cGood Functioning vs. Low Functioning.

Table 2. Variables potentially affecting the three functional profiles.

Characteristics	Good Functioning (n=926) No./mean	%/SD	Intermediate Functioning (n=1436) No./mean	%/SD	Low Functioning (n=661) No./mean	%/SD	F / χ^2	p-value
Age	38.5*	13.88	32.90 ^d	11.48	31.35 ^e	10.81	84.66	<.001
Sex, female n (%) ^a	737	79.8	1212	84.8	594	91.1	37.24	<.001
Work Situation							126.52	<.001
Employed	772	85.1	1219	85.1	483	73.1		
Unemployed	82	8.9	173	12.1	165	25		
Retired/Retired on disability	70	7.6	41	2.9	13	2		
Income (BRL) ^b							244.21	<.001
<708,19 - 2.965,69	207	22.4	498	34.7	369	55.8		
> 2.965,69 - 10.386,52	448	48.4	706	49.2	241	36.5		
> 10.386,52	271	29.3	232	16.2	51	7.7		
Occupation							29.00	<.001
Essential	312	33.7	406	28.3	141	21.3		
Non-essential	614	66.3	1030	71.7	520	78.7		
Previous psychiatric illness							146.44	<.001
Yes	264	28.5	598	41.6	389	31.1		
No	662	71.5	838	58.4	272	41.1		
Marital status							43.23	<.001
Married	488	52.7	594	41.4	249	37.7		
Single	438	47.3	842	58.6	412	62.3		
Education							70.16	<.001
Undergraduate	333	36	593	41.3	374	56.6		
Graduate/Postgraduate	593	64	843	58.7	287	43.4		
Impact of Event (IES-R)							640.46	<.001
Negative	839	90.6	977	68	198	30		
Positive	87	9.4	459	32	463	70		
Depression (PROMIS)							701.68	<.001
Moderate	322	34.8	1086	75.6	621	93.9		
Anxiety (PROMIS)							480.27	<.001
Moderate	581	62.7	1311	91.3	653	98.8		

Note: Different symbols mean difference between functioning conditions.

^a N=3005.

^b IBRL= 0.574

4.4. Variables Potentially Affecting the Overall FAST Score

As shown in Table 2, concerning sociodemographic variables, the one-way ANOVA with Tukey HSD post-hoc test and χ^2 showed differences among the three functioning groups in all characteristics.

As shown in Table 2, concerning sociodemographic variables, the one-way ANOVA with Tukey HSD post-hoc test and χ^2 showed differences among the three functioning groups in all characteristics.

In the multinomial regression analysis, the model showed a good fit to the data (Deviance: $\chi^2 = 4756,724$, $df = 5968$, $p = 1.00$; Nagelkerke's $R^2 = 0.455$) and was significant to account for variance in the cluster functioning (Model $\chi^2 (56) = 1529,46$, $p < 0.001$). The multinomial regression showed that higher scores in PROMIS depression (OR = 1.21(1.18 – 1.24) 95%CI, $p < 0.001$) and PROMIS anxiety (OR = 1.05, 95%CI (1.03 to 1.08), $p < 0.001$) significantly predicted whether the responder belongs to the low functioning cluster or the good functioning cluster, with a high relative risk belonging to the low functioning cluster. Furthermore, lower (OR = 3.42, 95%CI (2.18 to 5.35), $p < 0.001$) and middle (OR = 1.65, 95%CI (1.09 to 2.49), $p < 0.01$) household income significantly predicted whether the responder belongs to the low functioning cluster or the good functioning cluster, with higher odds belonging to the low functioning cluster. On the other hand, no previous history of psychiatric disorder (OR = 0.53, 95%CI (0.40 to 0.34), $p < 0.001$) and no symptoms of post-traumatic stress disorder (OR: 0.24, 95%CI (0.18 to 0.34), $p < 0.001$) significantly predicted whether the responder belongs to the low or the good functioning cluster, with a lower odd belonging to the low functioning cluster.

Furthermore, multinomial regression showed that younger age (OR: 0.99, 95%CI (0.98 to 1.00), $p < 0.05$), higher PROMIS depression (OR: 1.09, 95%CI (1.08 to 1.11), $p < 0.001$), and PROMIS anxiety scores (OR: 1.04, 95%CI (1.01 to 1.05), $p < 0.001$) significantly predicted whether the responder belongs to the intermediate or the good functioning cluster, with a high relative risk belonging to the intermediate functioning cluster. Moreover, lower (OR: 1.87, 95%CI 1.39 to 2.51, $p < 0.001$) and middle household income (OR: 1.40, 95%CI 1.10 to 1.79, $p < 0.01$) significantly predicted whether the responder belongs to the intermediate functioning cluster or the good functioning cluster, with higher odds belonging to the intermediate functioning cluster. However, lower education (OR = 0.72, 95%CI 0.58 to 0.89, $p < 0.01$) and no symptoms of post-traumatic stress disorder (OR = 0.58, 95%CI 0.44 to 0.77, $p < 0.001$) significantly predicted whether the responder belongs to the intermediate or the good functioning cluster, with a lower odds belonging to the intermediate functioning cluster.

5. DISCUSSION

The present study evaluated the psychometric properties of the online self-reported FAST scale and psychosocial implications of COVID-19 in a general population during the first peak of SARS-CoV-2 transmission in Brazil. The results showed that the online FAST scale retained the same six

domains as the original version, and the items had high internal consistency, with Cronbach's alpha above 0.9. In addition, the online version showed evidence of discriminant validity by differentiating three categories of psychosocial functioning in the sample and related variables with an overall FAST score. Our results showed the applicability of the digital FAST scale to assess functioning in the general population and revealed the main predictors of functional impairment during the first wave of SARS-CoV-2 contagion.

The ongoing pandemic imposes barriers for researchers worldwide, with many studies using online surveys to assess mental health in the general population [6, 30]. However, many instruments used in online studies have not been previously validated in the digital format and do not always reproduce the psychometric properties of original versions of the scales [31 - 33]. For instance, the assessment of anxiety in patients with panic disorder by the internet-based Beck Anxiety Inventory (BDI) questionnaire showed a significant difference in mean scores, with lower scores observed in the internet version compared to the original version of the scale [30]. In the present study, the internal consistency and reliability were found to be similar to a study that assessed the psychometric properties of the older version in bipolar patients and healthy controls by establishing a six-factor internal structure and a Cronbach's alpha of 0.9109 [13]. Furthermore, the psychometric properties of FAST in the present study were similar to the findings reported in a study on a subsample of adults with Autism Spectrum Disorder (ASD) or a sample involving patients in first-episode psychosis that also showed a six-factor structure and a Cronbach's alpha of 0.91 and 0.88, respectively [14, 18]. In addition, analysis of FAST reliability and factorial structure performed in patients with first-episode psychosis and healthy controls showed adequate reliability (Cronbach's alpha of 0.882) and a six-factor structure, suggesting that the FAST scale is applicable to a range of health conditions. However, mean scores of a high-functioning cluster were slightly higher than the cut-off observed in the older version [13], suggesting that misclassification of functional status would occur in the web survey if we applied a cut-off of the older version of FAST. Therefore, studies assessing the reliability and validity of new digital instruments, even those shifting from pen and paper format, are now required as many web surveys are in progress. In addition to reliability, the self-reported online FAST discriminated subjects into three clusters of functioning as follows: (I) good functioning group, representing one-third of individuals that experienced satisfactory functioning in distinct life domains, (II) intermediate functioning group, representing almost 50% of individuals with mild deficits in domains of functioning, and (III) low functioning group representing around 20% of individuals that experienced global and significant impairment in all domains of functioning. Some variables, namely age, income, mental health, and history of psychiatric disorder, significantly explain these functioning clusters. Responders in the LF cluster experienced greater anxiety and depressive symptoms as they self-reported more past events related to psychiatric disorders than other groups; therefore, both factors correlated to poor outcomes. Indeed, psychiatric symptoms and cognitive deficits have been traditionally associated with a

higher overall FAST score or poor functioning in clinical samples [13, 34].

The LF cluster also had lower socioeconomic levels and more financial difficulties than the other two clusters. Indeed, impairment in occupational functioning and financial issues might be a consequence of COVID-19 and pandemic preventive measures since there was an increase in unemployment rates, thus leading to more financial strain [35, 36]. The economic impact of the pandemic may aggravate the condition of more vulnerable individuals that, unfortunately, represent a huge part of the population in developing countries, like Brazil. Finally, younger people that were more prevalent in LF than other groups also reported more negative effects of COVID-19 than older subjects. Probably, pre-pandemic distress, such as educational, professional, or social difficulties typically experienced by young adults, compounded by lifestyle disruptions and feelings of hopelessness during the pandemic, may have contributed to these findings [37]. Additionally, they are more vulnerable to stressful situations because of their inexperience and lack of adaptive mechanisms. Together, these findings support that the digital FAST scale was sensitive to detect differences in functioning in a large sample of discriminating individuals by clinical symptoms and demographic characteristics, highlighting the potential utility of this scale in clinical and research settings.

One of the strengths of this instrument is that it is being validated in several cultures in either clinical samples or the general population. The self-reported online version showed strong psychometric properties, which are quite similar to its original version. The FAST may contribute to a multidimensional assessment of functioning with the advantage of being one of the very few validated scales in a digital format in a large community sample.

Nevertheless, some important limitations should be mentioned. Firstly, we used an online survey with a convenience sample method. Secondly, the instruments used to assess mental health were self-reported and might not characterize mental health status with the accuracy of structured clinical interviews. Also, although the digital version of the FAST showed satisfactory psychometric properties, we could not compare scores between different formats of the instrument to assess intra-class coefficients and mean scores; both measures are used to evaluate the degree of reliability and equivalence of the same instrument delivered using different formats [38]. Finally, it is noteworthy that online survey methods may have an issue of biased sampling toward people with good internet literacy and access.

CONCLUSION

The digital version of the FAST showed strong psychometric properties in the general population sample, indicating that the instrument measures a multidimensional construct of functioning, encouraging its use by researchers and clinicians in their practice. Moreover, these findings would help to better understand the psychosocial implications of the pandemic and the importance of planning specific interventions to rehabilitate the affected group. Considering the previous reports [6], it can be concluded that mental health problems and poor psychosocial functioning may be a mark left by this pandemic.

LIST OF ABBREVIATIONS

WHO	=	World Health Organization
COVID-19	=	Severe Acute Respiratory Syndrome Coronavirus
FAST	=	Functioning Assessment Short Test
IES-R	=	Impact of Event Scale-Revised
PROMIS	=	Patient-Reported Outcomes Measurement Information System
CFA	=	Confirmatory Factor Analysis
KMO	=	Kaiser-Meyer-Olkin
PAM	=	Partition Around Medoids

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

All procedures performed in studies involving human participants were in accordance with the ethical standards of the local ethics committee (Hospital de Clínicas de Porto Alegre, protocol number 2020-0162).

HUMAN AND ANIMAL RIGHTS

No animals were used in this research. All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008.

CONSENT FOR PUBLICATION

Written informed consent was obtained from each participant prior to the study for publication of this research.

STANDARDS OF REPORTING

STROBE guideline has been followed.

AVAILABILITY OF DATA AND MATERIALS

Data supporting the findings of the article is not publicly available as a result of the privacy policies of the health facilities involved in the study, but it can be provided by the corresponding author [A.R.R.] upon reasonable request.

FUNDING

This study was financed by FIPE-HCPA GPPG no. 2020-0162. This study was partially funded by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Coordination for the Improvement of Higher Education Personnel; CAPES) - Brasil - Finance Code 001.

CONFLICT OF INTEREST

E.V. has received grants and served as a consultant, advisor or CME speaker unrelated to this work for the following entities: AB-Biotics, Abbvie, Angelini, Daiippon Sumitomo Pharma, Ferrer, GH Research, Gedeon Richter, Janssen, Lundbeck, Otsuka, Sage, Sanofi-Aventis, Sunovion, and Takeda. The other authors do not declare any conflict of interest related to this manuscript.

ACKNOWLEDGMENTS

Adriane R Rosa would like to thank the CNPq Conselho Nacional de Desenvolvimento Científico e Tecnológico - Brasil process 302382/2019-4. Jéferson F Goularte would like to acknowledge Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES). Silvia D Serafim would like to thank CNPq.

REFERENCES

- [1] World Health Organization. Coronavirus Disease (COVID-19) outbreak 2020. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> [cited 2021 Nov 13].
- [2] Kola L, Kohrt BA, Hanlon C, et al. COVID-19 mental health impact and responses in low-income and middle-income countries: Reimagining global mental health. *Lancet Psychiatry* 2021; 8(6): 535-50. [[http://dx.doi.org/10.1016/S2215-0366\(21\)00025-0](http://dx.doi.org/10.1016/S2215-0366(21)00025-0)] [PMID: 33639109]
- [3] Afifi RA, Novak N, Gilbert PA, et al. 'Most at risk' for COVID-19? The imperative to expand the definition from biological to social factors for equity. *Prev Med* 2020; 139: 106229. [<http://dx.doi.org/10.1016/j.ypmed.2020.106229>] [PMID: 32763263]
- [4] Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *JAMA* 2020; 323(21): 2133-4. [<http://dx.doi.org/10.1001/jama.2020.5893>] [PMID: 32259193]
- [5] Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *Lancet* 2020; 395(10227): 912-20. [[http://dx.doi.org/10.1016/S0140-6736\(20\)30460-8](http://dx.doi.org/10.1016/S0140-6736(20)30460-8)] [PMID: 32112714]
- [6] Goularte JF, Serafim SD, Colombo R, Hogg B, Caldieraro MA, Rosa AR. COVID-19 and mental health in Brazil: Psychiatric symptoms in the general population. *J Psychiatr Res* 2021; 132: 32-7. [<http://dx.doi.org/10.1016/j.jpsychires.2020.09.021>] [PMID: 33038563]
- [7] Rossi R, Socci V, Talevi D, et al. COVID-19 pandemic and lockdown measures impact on mental health among the general population in Italy. *Front Psychiatry* 2020; 11: 790. [<http://dx.doi.org/10.3389/fpsy.2020.00790>]
- [8] Vahia IV, Blazer DG, Smith GS, et al. COVID-19, mental health and aging: A need for new knowledge to bridge science and service. *Am J Geriatr Psychiatry* 2020; 28(7): 695-7. [<http://dx.doi.org/10.1016/j.jagp.2020.03.007>] [PMID: 32278745]
- [9] Chi X, Liang K, Chen ST, et al. Mental health problems among Chinese adolescents during the COVID-19: The importance of nutrition and physical activity. *Int J Clin Health Psychol* 2021; 21(3): 100218. [<http://dx.doi.org/10.1016/j.ijchp.2020.100218>] [PMID: 33391373]
- [10] Lima F, Da Ponte RFD, Bücken J, et al. Identifying cognitive subgroups in bipolar disorder: A cluster analysis. *J Affect Disord* 2019; 246: 252-61. [<http://dx.doi.org/10.1016/j.jad.2018.12.044>] [PMID: 30590288]
- [11] Michalak EE, Murray G. A clinician's guide to psychosocial functioning and quality of life in bipolar disorder. In: Young A, Ferrier N, Michalak EE, Eds. *Practical Management of Bipolar Disorder* 2010; 163-74.
- [12] Öngür D, Perlis R, Goff D. Psychiatry and COVID-19. *JAMA* 2020; 324(12): 1149-50. [<http://dx.doi.org/10.1001/jama.2020.14294>] [PMID: 32960247]
- [13] Rosa AR, Sánchez MJ, Martínez AA, et al. Validity and reliability of the Functioning Assessment Short Test (FAST) in bipolar disorder. *Clin Pract Epidemiol Ment Health* 2007; 3: 5. [<http://dx.doi.org/10.1186/1745-0179-3-5>]
- [14] Gisbert GL, Lugo MJ, Ramos IS, et al. Functioning Assessment Short Test (FAST): Validity and reliability in adults with autism spectrum disorder. *BMC Psychiatry* 2021; 21(1): 321. [<http://dx.doi.org/10.1186/s12888-021-03330-y>]
- [15] Zhang Y, Long X, Ma X, et al. Psychometric properties of the Chinese version of the Functioning Assessment Short Test (FAST) in bipolar disorder. *J Affect Disord* 2018; 238: 156-60. [<http://dx.doi.org/10.1016/j.jad.2018.05.019>] [PMID: 29883937]
- [16] Rotger S, Richarte V, Nogueira M, et al. Functioning Assessment Short Test (FAST): Validity and reliability in adults with attention-deficit/hyperactivity disorder. *Eur Arch Psychiatry Clin Neurosci* 2014; 264(8): 719-27. [<http://dx.doi.org/10.1007/s00406-014-0501-0>] [PMID: 24710954]
- [17] Ruiz A, Pousa E, Duñó R, Crosas J, Cuppa S, García C. Adaptation to Spanish of the assessment Scale Of Unawareness Of Mental Disorder (SUMD). *Actas Esp Psiquiatr* 2008; 36(2): 111-1198. [Spanish adaptation of the Scale to Assess Unawareness of Mental Disorder (SUMD)]. [PMID: 18365791]
- [18] Amoretti S, Mezquida G, Rosa AR, et al. The Functioning Assessment Short Test (FAST) applied to first-episode psychosis: Psychometric properties and severity thresholds. *Eur Neuropsychopharmacol* 2021; 47: 98-111. [<http://dx.doi.org/10.1016/j.euroneuro.2021.02.007>] [PMID: 33658165]
- [19] Cacilhas AA, Da Silva MPV, Ceresér KM, et al. Validity of a Short Functioning Test (FAST) in Brazilian outpatients with bipolar disorder. *Value Health* 2009; 12(4): 624-7. [<http://dx.doi.org/10.1111/j.1524-4733.2008.00481.x>] [PMID: 19900260]
- [20] Barbato A, Bossini L, Calugi S, et al. Validation of the Italian version of the Functioning Assessment Short Test (FAST) for bipolar disorder. *Epidemiol Psychiatr Sci* 2013; 22(2): 187-94. [<http://dx.doi.org/10.1017/S2045796012000522>] [PMID: 23089135]
- [21] Akers N, Lobban F, Hilton C, Panagaki K, Jones SH. Measuring social and occupational functioning of people with bipolar disorder: A systematic review. *Clin Psychol Rev* 2019; 74: 101782. [<http://dx.doi.org/10.1016/j.cpr.2019.101782>] [PMID: 31751878]
- [22] Riegler C, Wiedmann S, Rücker V, et al. A self-administered version of the functioning assessment short test for use in population-based studies: A pilot study. *Clin Pract Epidemiol Ment Health* 2020; 16: 192-203. [<http://dx.doi.org/10.2174/1745017902016010192>]
- [23] Orhan M, Kortén N, Kupka R, et al. Reliability and validity of the functioning assessment short test for older adults with bipolar disorder (FAST-O). *Int J Bipolar Disord* 2020; 8(1): 28. [<http://dx.doi.org/10.1186/s40345-020-00193-2>]
- [24] Van Ameringen M, Turna J, Khalesi Z, Pullia K, Patterson B. There is an app for that! The current state of mobile applications (apps) for DSM-5 obsessive-compulsive disorder, posttraumatic stress disorder, anxiety and mood disorders. *Depress Anxiety* 2017; 34(6): 526-39. [<http://dx.doi.org/10.1002/da.22657>] [PMID: 28569409]
- [25] Van Ballegooijen W, Riper H, Cuijpers P, Van Oppen P, Smit JH. Validation of online psychometric instruments for common mental health disorders: A systematic review. *BMC Psychiatry* 2016; 16: 45. [<http://dx.doi.org/10.1186/s12888-016-0735-7>]
- [26] Caiuby AVS, Lacerda SS, Quintana MI, Torii TS, Andreoli SB. Cross-cultural adaptation of the Brazilian version of the Event Impact Scale - Revised (IES-R). *Cad Saude Publica* 2012; 28(3): 597-603. [<http://dx.doi.org/10.1590/S0102-311X2012000300019>] [PMID: 22415191]
- [27] Taherdoost H, Sahibuddin S, Jalaliyoon N. Exploratory factor analysis; concepts and theory. *Adv Appl Pure Math* 2014; 27: 375-82.
- [28] Cohen J. *Statistical Power for the Behavioral Sciences*. New York: Academic Press 1977.
- [29] Schubert E, Rousseeuw PJ. Faster k-medoids clustering: Improving the PAM, CLARA, and CLARANS algorithms. *International conference on similarity search and applications*. Cham:Springer. 2019; pp. 171-87. [http://dx.doi.org/10.1007/978-3-030-32047-8_16]
- [30] Xiong J, Lipsitz O, Nasri F, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord* 2020; 277: 55-64. [<http://dx.doi.org/10.1016/j.jad.2020.08.001>] [PMID: 32799105]
- [31] Carlbring P, Brunt S, Bohman S, et al. Internet vs. paper and pencil administration of questionnaires commonly used in panic/agoraphobia research. *Comput Human Behav* 2007; 23(3): 1421-34. [<http://dx.doi.org/10.1016/j.chb.2005.05.002>]
- [32] Maguire Á, Martin J, Jarke H, Ruggeri K. Getting closer? Differences remain in neuropsychological assessments converted to mobile devices. *Psychol Serv* 2019; 16(2): 221-6. [<http://dx.doi.org/10.1037/ser0000307>] [PMID: 30407051]
- [33] Ruggeri K, Maguire Á, Andrews JL, Martin E, Menon S. Are we there yet? Exploring the impact of translating cognitive tests for dementia using mobile technology in an aging population. *Front Aging Neurosci*

- 2016; 8: 21.
[<http://dx.doi.org/10.3389/fnagi.2016.00021>]
- [34] Grande I, Goikolea JM, De Dios C, *et al.* Occupational disability in bipolar disorder: Analysis of Predictors Of Being On Severe Disablement Benefit (PREBIS study data). *Acta Psychiatr Scand* 2013; 127(5): 403-11.
[<http://dx.doi.org/10.1111/acps.12003>] [PMID: 22924855]
- [35] Godinic D, Obrenovic B, Khudaykulov A. Effects of economic uncertainty on mental health in the COVID-19 pandemic context: Social identity disturbance, job uncertainty and psychological well-being model. *Int J Innov Econ Dev* 2020; 6(1): 61-74.
[<http://dx.doi.org/10.18775/ijied.1849-7551-7020.2015.61.2005>]
- [36] Moretti A, Menna F, Alicino M, Paoletta M, Liguori S, Iolascon G. Characterization of home working population during COVID-19 emergency: A cross-sectional analysis. *Int J Environ Res Public Health* 2020; 17(17): 6284.
[<http://dx.doi.org/10.3390/ijerph17176284>]
- [37] Shanahan L, Steinhoff A, Bechtiger L, *et al.* Emotional distress in young adults during the COVID-19 pandemic: Evidence of risk and resilience from a longitudinal cohort study. *Psychol Med* 2022; 52(5): 824-33.
[<http://dx.doi.org/10.1017/S003329172000241X>] [PMID: 32571438]
- [38] Alfonsson S, Maathz P, Hursti T. Interformat reliability of digital psychiatric self-report questionnaires: A systematic review. *J Med Internet Res* 2014; 16(12): e268.
[<http://dx.doi.org/10.2196/jmir.3395>]

© 2022 Serafim *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.